

A review of the stratigraphy of Eastern Paratethys (Oligocene–Holocene)

R.W. JONES

BP Exploration, Uxbridge One, 1 Harefield Rd., Uxbridge, Middlesex, UB8 1PD, U.K.

M.D. SIMMONS

Department of Geology and Petroleum Geology, University of Aberdeen, Meston Building, King's College, Aberdeen, AB9 2UE, U.K.

CONTENTS

Introduction	26
Absolute Chronostratigraphy	26
Micropalaeontological Biostratigraphy and Palaeoenvironmental Interpretation	26
Biostratigraphy	26
Palaeoenvironmental Interpretation	28
<i>Non-Marine Environments</i>	28
<i>Quasi-Marine and Marine Environments</i>	28
Climatostratigraphy	29
Magnetostratigraphy	29
Oxygen Isotope Stratigraphy	29
Sequence Stratigraphy and Palaeogeography	29
Introduction	29
Maykopian	33
Tarkhanian to Konkian	34
Tarkhanian	35
Chokrakian	35
Karaganian	35
Konkian	36
Sarmatian	36
Maeotian	38
Pontian	38
Kimmerian	39
Akchagylia to Khvalynian (Caspian Sea)	40
Akchagylia	40
Apsheronian	42
Bakunian	42
Khazarian, Girkan and Khvalynian	42
Kuyalnikian to Neoeuxinian (Black Sea)	42
Kuyalnikian	42
Gurian	43
Chaudian	43
Uzunlarian, Karangatian and Neoeuxinian	43
Acknowledgements	43
References	43

SYNOPSIS. All available data pertaining to the regional stratigraphy of Eastern (Ponto-Caspian) Paratethys, much of it in sources not freely available in the west, is reviewed. Particular emphasis is placed on the South Caspian. Where possible, regional datums are calibrated against global standards. An attempt is made to place the regional stratigraphy in a (global) sequence stratigraphic framework for the first time. Palaeogeographic reconstructions are given for selected time-slices.

INTRODUCTION

The Tethyan Ocean began to close in the Eocene as a result of plate collisions along the southern margin of the Eurasian Supercontinent that ultimately gave rise to the formation of the mountain chain extending from the Alps in the west to the Himalayas in the east. The initial response to these plate collisions was the formation of a suite of east-west trending sedimentary basins extending from Austro-Hungary in the west to Central Asia in the east, collectively constituting the intracontinental Paratethyan Sea (Fig. 1). Subsequent tectonic uplift (enhanced by eustatic shallowing) through the Mio-Pliocene led to widespread marginal- to non- marine sedimentation. Ultimate severance of connections to the world's oceans led to the evolution of largely endemic faunas and floras (in particular in Eastern Paratethys, which was more isolated than Central Paratethys). This renders stratigraphic correlation between established Mediterranean and Paratethyan stages extremely difficult. The problem is locally compounded by confusion between chronostratigraphic and lithostratigraphic nomenclature.

In this paper we review all available data, much of it in sources not freely available in the west, on the regional stratigraphy of Eastern Paratethys (Fig. 1). We place particular emphasis on the South Caspian, an area in which the western oil industry is showing a growing interest, and one with which we, through our industrial work and our academic contacts in the Former Soviet Union, are particularly familiar.

We give an indication of the palaeontology of each regional stage. For the sake of brevity and because of their stratigraphic utility, we concentrate on various groups of microfossils, though we acknowledge that macrofossils, especially molluscs, also have stratigraphic value (see, for instance, Ali-Zade (1954), Azizbekov (1972), Ali-Zade *et al.* (1986), (Azerbaijan); Lupov *et al.* (1972) (Turkmenia); Andreescu (1981) (Dacic Basin); and Ozsayar (1985) and Taner (1985) (Turkey)). A forthcoming paper (Simmons *et al.* in press) will document in detail the micropalaeontological (including nannopalaeontological and palynological) and macropalaeontological zonation of the

Neogene to Pleistocene sediments of Azerbaijan.

We attempt to place the regional stratigraphy in a global framework by calibrating biostratigraphic and magnetostratigraphic datums against Central Paratethyan and Mediterranean standards, and by suggesting possible calibrations between regional sequence boundaries and flooding surfaces, and the global sequence stratigraphic framework and eustatic sea-level curve of Haq *et al.* (1988).

ABSOLUTE CHRONOSTRATIGRAPHY

Notwithstanding the efforts of such authors as Steininger & Papp (1979), Chumakov *et al.* (1984, 1988, 1992a-b) and Vass (1985), there is no established comprehensive absolute chronostratigraphic time-scale for Eastern Paratethys. Thus, in Eastern Paratethys, absolute chronostratigraphic dating is often only possible by calibration of regional stratigraphic datums against global standards. We have attempted to calibrate regional datums against the Haq *et al.* (1988) timescale, which is the most up-to-date timescale that conveniently integrates bio-, magneto- and sequence- stratigraphic data. The confidence with which this sort of calibration can be made varies considerably with stratigraphic interval (see below).

MICROPALAEONTOLOGICAL BIOSTRATIGRAPHY AND PALAEOENVIRONMENTAL INTERPRETATION

Biostratigraphy

Those groups of planktonic organisms traditionally used in the biostratigraphic zonation of the Cenozoic (planktonic foraminifera and calcareous nannoplankton) are restricted in their development in Paratethys because of the isolated




Fig. 1 Geological sketch map of the Paratethyan Basin in the Oligocene. Modified after Steininger & Papp (1979). C = Central Paratethys; E = Eastern Paratethys. Eastern Paratethys can be considered as including the Pontian (Black Sea) Basin and the Caspian Basin. Throughout most of the Miocene, the eastern limit of the Paratethyan Basin was probably the Caspian Basin, but in the Late Pliocene at least it was probably further east once more.

geological evolution of the region. Only locally or periodically (as in the Maykopian, Maeotian and Kuyalnikian/Akchagylian (see section below on sequence stratigraphy)) are they sufficiently well developed to enable ties to global biostratigraphic zonation schemes (Blow (1969) for planktonic foraminifera; Martini (1971) for calcareous nannoplankton).

Biostratigraphic zonation in Eastern Paratethys relies largely on facies-dependent benthonic foraminifera and, especially in the marginal- to non-marine environments of the Mio-Pliocene, benthonic ostracods and terrestrially-derived pollen and spores (see section below on sequence stratigraphy). Important ostracod references include those of Livental (1929), Sveier (1949), Agalarova (1956, 1967), Suzin (1956), Agalarova *et al.* (1961), Mandelstam *et al.* (1962), Faridi (1964), Imnadze (1964,

1974), Sheydayeva-Kuliyeva (1966), Rozyeva (1971), Gramann (1971), Karmishina (1975), Vekua (1975), Krstic (1976), Olteanu (1978), Imnadze & Karmishina (1980), de Deckker (1981), Jiricek (1984), Mamedova (1984, 1985, 1988), Dzhaneldidze *et al.* (1985), Aliyulla *et al.* (1985) and Yassini (1986). Important pollen and spore and associated palynomorph references include those of Ramishvili (1969), Dzhabarova (1973, 1978, 1980), Grichuk (1973, 1984), Wall & Dale (1973), Ananova (1974), Shikmus *et al.* (1977), Koreneva & Kartashova (1978), Traverse (1978), Shchekina (1979), Abramova (1982, 1985), Yakhimovich *et al.* (1983), Grichuk *et al.* (1984), Khotinskiy (1984), Mamedov & Rabotina (1984a–b), Shatilova (1984), Ananova *et al.* (1985), Ivanova (1985), Sirenko & Turlo (1986), Bludorova *et al.* (1987), Naidina (1988, 1990a–b,




Fig. 2 Vegetation belts in the former Soviet Union. After Knystautas (1987). A.D. = Amu Darya; S.D. = Syr Darya.

1991a–c) and Malaeva & Kulikov (1991).

Other locally stratigraphically useful fossil groups include otoliths (Brzobohaty, 1983), Problematica (*Bolboforma* (Szczuchura, 1985; Spiegler & Rögl, 1992)), siliceous microfossils (diatoms (Shishova, 1955; Ushakova & Ushko, 1971; Gasanova, 1965; Rasulov, 1986), radiolarians (Slama, 1983), silicoflagellates (Dumitrica, 1985) and sponge spicules (Riha, 1983)), and, in non-marine environments, vertebrate remains (*Camelopardis*, *Felis*, *Gazella*, *Hipparion*, *Hyaena*, *Mastodon*, *Mesopithecus*, *Rhinoceros*, etc.) (Kretzoi, 1985; Steininger, Rabeder & Rögl, 1985; Bernor *et al.*, 1987, 1993; Lindsay *et al.*, 1989; Rögl *et al.*, 1993) and charophytes (Rögl *et al.*, 1993).

Ali-Zade *et al.* (1994a, 1994b, 1995, in press), Reynolds *et al.* (in press) and Simmons *et al.* (in press) give further details of the Neogene biostratigraphy of Eastern Azerbaijan.

Palaeoenvironmental Interpretation

Non-Marine Environments. Non-marine environments are characterised by fresh-water ostracods such as *Aglaioecypris*, *Candona*, *Candonella*, *Cyclocypris*, *Cypria*, *Eucypris*, *Ilyocypris*, *Pseudostenocypris* and *Zonocypris*, and terrestrially-derived pollen and spores. Pennate diatoms, fresh-water gastropods and terrestrial vertebrate remains may also be found.

Palaeoclimate can be inferred from the distribution of vegetation types as inferred from pollen and spores. At the present-day, the distribution of vegetation types is determined chiefly by climatic factors (temperature (latitude, altitude), aridity). Thus, for instance, birches characterise the cold 'forest-tundra' of the extreme north, diverse coniferous and deciduous types the 'taiga' of the central area, and grasses and shrubs the arid treeless 'steppe' and semi-desert to the extreme south (Figs 2–3).

Quasi-Marine and Marine Environments. Deposition in oligo- to meso- haline (hereafter 'quasi-marine' (brackish, reduced salinity)) environments prevailed in the Paratethyan Basin (especially in the Caspian) throughout much of its geological evolution because of its restricted connection to the open ocean (see above). However, water depths and sedimentary regimes may have been similar to those of the normal marine realm, and, moreover, deposition under normal or near-normal marine conditions did take place at times (e.g., Maykopian and Akchaglyan). Palaeosalinity can be inferred from diatoms (e.g., Ushakova & Ushko, 1971; Schrader, 1979) or from foraminifera and ostracods ranging through to the Recent (see below). Palaeosalinity and/or palaeotemperature curves are presented by Semenenko (1979), Chepalyga (1985) and Demarcq (1985).

Quasi-marine environments in the Black Sea and Caspian Sea are characterised by the benthonic foraminiferal genus *Florilus*, and some species of the genera *Ammobaculites*, *Ammoscalaria*, *Ammonia* and *Elphidium* (salinity tolerance range 1–5 parts per thousand (ppt)), and *Miliammina*, *Haynesina* and *Rosalina* and some species of *Nonion s.l.* and *Quinqueloculina* (1–26ppt) (Macarovič & Cehan-Ionesi, 1962; Tufescu, 1968, 1973; Murray, 1973, 1991; Gheorghian, 1974; Yassini & Ghahreman, 1977; Yanko, 1990b), the ostracod genera *Cyprideis* (2–14ppt), *Maetocythere* (4–14ppt), *Loxococoncha* (5–14ppt), *Bakunella*, *Caspiolla* and *Cytherissa* (11–13/14ppt) and *Graviacypris* (12–13ppt) (Gofman, 1966; Yassini, 1986; Boomer, 1993a), and the calcareous nannofossil genus *Emiliania* (11ppt) (Bukry, 1974).

Normal or near-normal marine environments are

characterised by the benthonic foraminiferal genera *Discorbis*, *Textularia*, *Bolivina*, *Bulimina*, *Brizalina*, *Cibicides*, *Gavelinopsis* and *Trifarina* and some species of the genera *Ammonia*, *Nonion s.l.* and *Quinqueloculina* (salinity tolerance range 11–26ppt) (Macarovič & Cehan-Ionesi, 1962; Tufescu, 1968, 1973; Murray, 1973, 1991; Yassini & Ghahreman, 1977; Yanko, 1990b).

CLIMATOSTRATIGRAPHY

Zubakov & Borzenkova (1990) defined a series of climatostratigraphic units called 'climathems' (some conceptual, some stratotypified (and with representative pollen spectra documented)) which they used in the regional correlation of Eastern Paratethys (see also Zubakov, 1993). Of these, 'superclimathems' (SCTs), with an average duration of 200,000 years, are the most useful. SCTs are correlated with half the 370,000–425,000-year cycle of orbital eccentricity, and reflect changes in climate (alternating between 'cryo-' and 'thermo-' meric (cool and warm respectively)).

Zubakov & Borzenkova interpreted pollen spectra dominated by steppe and semi-desert vegetation as being of 'warm' aspect (whereas, in fact, they are more characteristic of aridity than high temperature) and those dominated by forest vegetation as being of 'cool' aspect (whereas they are in fact more characteristic of humidity than of low temperature). In the Caspian, they found the former to characterise regressions and the latter to characterise transgressions, and therefore correlated regressions with 'warm' phases (interglacials) and transgressions with 'cool' phases (glacials) (Fig. 4A). Regression during interglacials is possible if sediment supply and subsidence are in equilibrium but evaporation exceeds precipitation and run-off. Transgression during glacials is possible if sediment supply (reduced by rivers freezing) fails to fill the accommodation space created by subsidence. Note in this context that the level of the Caspian has fallen by some 5m since records were first taken in the 1760's. However, it also appears to have been rising over the last fifty years (currently at a rate of 20cms/year in the South Caspian). Historical records are probably unreliable in a geological context because of the influence of man on the environment. This is particularly true in the case of the Aral Sea, whose level has fallen and whose salinity has risen drastically since the 1960's owing to abstraction of the headwaters of the feeder rivers (the Amu Darya and Syr Darya) to irrigate the cotton fields of Uzbekistan (see, for instance, Boomer, 1993a–b). Incidentally, as a result of this catastrophic ecological change, eleven species of fresh-water ostracod known to have been living in the Aral Sea thirty years ago no longer live there. Only the quasi-marine species *Cyprideis torosa* lives there today.

The evidence for regressions during interglacials and transgressions during glacials appears somewhat equivocal. An equally strong case, and one more in keeping with *a priori* expectation from experience in other parts of the world, can be made for correlating regressions with glacials and transgressions with interglacials (Fig. 4B). One key observation in support of this case is the apparent correlation of the major transgressions not only with warm phases (see, for instance, Skallbdyna, 1985), but also with global transgressions (see, for instance, Haq *et al.*, 1988). Pollen spectra of 'arid' aspect in glacial sediments and of 'humid' aspect in interglacial sediments are explicable in terms of, respectively, contractions and expansions of the forest belt




Fig. 3 Distribution of principal vegetation types in relation to vegetation belts. Compiled from various sources. Bar width provides a measure of abundance.

(the former in response to permafrost development). This is also indicated by palaeoclimatic reconstructions for the Late Valdai Glacial and Mikulino Interglacial (Grichuk, 1984; Savina & Khotinskiy, 1984; Velichko, 1984b). An alternative explanation envisages a time lag between the onset of glaciation and the response of vegetation (belts being displaced by up to 2000km.). Similar disequilibrium phenomena have been described from interstadial complexes in the United Kingdom.

MAGNETOSTRATIGRAPHY

Much magnetostratigraphic data is available from Eastern Paratethys (e.g., Zubakov & Kochegura (1971), Trubikhin (1977), Semenenko (1979), Semenenko & Pevzner (1979), Grishanov *et al.* (1983), Rögl & Steininger (1984), Steininger & Rögl (1984), Chepalyga (1985), Chepalyga *et al.* (1985), Iossofova (1985), Pevzner & Vangengeim (1985a, 1993), Senes (1985), Skalbldyna (1985), Vass (1985), Zubakov & Borzenkova (1990) and Trubikhin *et al.* (1991a–b)). Theoretically, this sort of data ought to enable a correlation between Eastern Paratethys and the rest of the world (which, as noted above, is difficult to do using the available biostratigraphic data). However, in practice the process is complicated by apparently inconsistent definition and usage of magnetostratigraphic units (polarity epochs). It is beyond the scope of this paper to address this problem in any more detail (instead, we simply quote the published magnetostratigraphic (polarity epoch) ranges for the various regional stages). It is nonetheless evident that the potential exists for a refined magnetostratigraphic subdivision of critical intervals using short-lived polarity reversal 'episodes' within the longer-term epochs.

OXYGEN ISOTOPE STRATIGRAPHY

Theoretically, the ages of the Plio-Pleistocene sediments of Eastern Paratethys are resolvable using oxygen isotope stratigraphic techniques. However, in practice, what data there is exists in widely disseminated form and is not particularly useful.

SEQUENCE STRATIGRAPHY AND PALAEOGEOGRAPHY


Introduction

No published sequence stratigraphic schemes exist for the Oligocene-Holocene of Paratethys, although relative sea-level changes and associated aspects of sequence stratigraphy are discussed by Rögl & Steininger (1983), Chepalyga (1985, 1991), Demarcq (1985), Krhovsky (1985), Nevesskaya *et al.* (1985), Pogacsas (1985), Pogacsas & Revesz (1985), Skalbldyna (1985), Andalibi (1991), Zubakov & Borzenkova (1990) and Klopovotskaya (1991).


This section attempts to place the regional stratigraphy of Eastern Paratethys in a (global) sequence stratigraphic framework. It is written in the form of a geological history. Stratigraphic (bio-, climato-, magneto- and sequence-stratigraphic) data are summarised on Figs 5–6. Palaeogeographic reconstructions for selected time-slices are given on Figs 7–12. These are based in part on previously published maps (Podobina *et al.*, 1956; Muratov, 1960; Sheydayeva-Kuliyeva, 1966; Ushakova & Ushko, 1971; Senes, 1973; Azizbekova, 1974; Senes & Marinescu, 1974; Lutttig &

A GLACIAL - TRANSGRESSION


- Rivers frozen. Extensive Permafrost.
- Sediment Supply < Subsidence.
- Expansion of 'Cool' Zone.

**INTERGLACIAL - REGRESSION**

- River Systems Reactivated.
- Evaporation > Precipitation / Runoff.
- Expansion of 'Warm' Zone.

**B GLACIAL - REGRESSION**

- Water locked up in Ice Sheet.
- Pollen Spectra of Arid Aspect.
- Contraction of Forest Zone

**INTERGLACIAL - TRANSGRESSION**

- Water released from melting Ice Sheet
- Expansion of Forest Zone




Fig. 4 Response to glaciation – alternative models for the Caspian. A: glacial transgression/interglacial regression. B: glacial regression/interglacial transgression. B better follows the palaeoclimatic reconstruction data given by Grichuk (1984) (see text).




Fig. 5 Stratigraphic summary (Eocene-Holocene). Chronostratigraphy, magnetostratigraphy, global sequence stratigraphy and calibration after Haq *et al.* (1988) (see also chronostratigraphic schemes of Chumakov *et al.*, 1984, 1988, 1992a-b and others). Sequence boundary ages in Ma. Biostratigraphy after Blow (1969) (planktonic foraminifera) (prefixed P and N) and Martini (1971) (calcareous nanoplankton) (prefixed NP and NN). Regional stratigraphy from this paper. Regional sequence stratigraphy modified after Chepalyga (1985). Cycles are regressive (mega) sequences. II-III are equivalent to the 'Eoparatethyan', IV to the 'Mesoparatethyan' and V-VI to the 'Neoparatethyan' of Nevesskaya *et al.* (1985). Curve shows extent of open marine connection (function of sea-level) as inferred from salinity data from palaeontological analyses (S‰ = salinity parts per thousand). Res. pot. = Resource potential (S = source, R = reservoir, C = caprock). The correlation of Eastern Paratethyan sequence stratigraphy with the global coastal onlap curve of Haq *et al.* (1988) is tentative. Where biostratigraphic control constrains the correlation this is indicated.

Steffens, 1976; Steininger, Rögl & Martini, 1976; Hsu, *in* Ross *et al.*, 1978; Rögl *et al.*, 1978; Steininger & Papp, 1979; Hsu, 1983; Baldi, 1984; Rögl & Steininger, 1983, 1984; Steininger & Rögl, 1984; Voronina & Popov, 1984; Iossofova, 1985; Popescu, 1985; Rusu, 1985; Steininger, Rabeder & Rögl, 1985; Steininger, Rögl & Nevesskaya, *in* Steininger, Senes, Kleeman & Rögl, 1985; Voicu, 1985; Bernor *et al.*, 1987; Nevesskaya *et al.*, 1987; Panakhi & Buare Mamadu Lamin, 1987; Veto, 1987; Mamedov, 1989; Olteanu, 1989; Adamia *et al.*, 1990; Dercourt *et al.*, 1990; Tchoumatchenko *et al.*, 1990; Kerimov *et al.*, 1991; Spiegler & Rögl, 1992; Pevzner & Vangengeim, 1993), and in part on previously unpublished maps.

It should be noted that the calibration against the global sequence stratigraphic framework and eustatic sea-level curve of Haq *et al.* (1988) is tentative. Fig. 5 demonstrates where biostratigraphic control exists in order to constrain the calibration. In the absence of such constraint, calibration is made by matching patterns of transgression and regression within a looser stratigraphic framework. The correlation between Eastern Paratethyan and global sequence stratigraphy and eustatic sea-level appears good, with all of the global eustatic sea-level trends finding their expression in Eastern Paratethys. It could be argued that this apparent correlation is entirely fortuitous. However, the stratigraphic signature of the mid-late Cenozoic appears remarkably consistent throughout the world, presumably because at this time it was an 'ice-house' world characterised by over-riding glacio-eustasy (Vail *et al.*, 1991). Indeed, it may be that not only third-order but also higher frequency sea-level oscillations are recognisable in areas characterised by a high sedimentation rate such as Eastern Paratethys (and the Gulf of Mexico (see, for instance, Beard *et al.*, 1982; Lamb *et al.*, 1987; Pacht *et al.*, 1990; Wornardt & Vail, 1990; see also Fig. 6)).

General features of Eastern Paratethyan stratigraphy have been discussed by, among others, Bogdanowicz (1947), Muratov (1960), Subbotina *et al.* (1960), Dzhaneldze (1970), Mamedova (1971, 1987), Stöcklin & Setudehnia (1971, 1972), Azizbekov (1972) (and authors cited therein), Lupov *et al.* (1972), Cicha *et al.* (1975), Jiricek (1975), Ross *et al.* (1978), Nikiforova & Dodonov (1980), Verisharin *et al.* (1982) (and authors cited therein), Alekseyev & Nikiforova (1984), Iossofova (1985), Popov & Voronina (1985), Semenenko & Lulieva (1985), Skalbdyna (1985), Volkova *et al.* (1985), Yakhemovich *et al.* (1985), Muratov & Nevesskaya (1986), Kereudren & Thibault (1987), Nigarov & Fedorov (1987), Benyamovskoy *et al.* (1988), Steininger *et al.* (1989), Yanko (1990a–b, 1991), Zubakov & Borzenkova (1990), Ghanbari (1991), Ali-Zade *et al.* (1994a, 1994b, 1995, *in press*), Jones (1996), Reynolds *et al.* (*in press*) and Simmons *et al.* (*in press*). Correlations within Eastern Paratethys and between Eastern and Central Paratethys have been discussed by Papp (1969), Chelidze (1973), Rögl, Steininger & Muller (1978), Paramonova *et al.* (1979), Semenenko (1979, 1984), Semenenko & Pevzner (1979), Steininger & Papp (1979), Steininger & Rögl (1979, 1984), Baldi (1980), Semenenko & Lulieva (1982), Rögl & Steininger (1983, 1984), Nevesskaya *et al.* (1984, 1985, 1987), Velichko (1984a), Yakhimovich, Bludorova, Zhidovinov *et al.* (1984), Chepalyga *et al.* (1985), Nevesskaya & Nosovsky (1985), Nosovsky (1985), Pevzner & Vangengeim (1985), Rögl (1985a), Senes (1985a–b), Senes & Steininger, *in* Steininger *et al.* (1985), Yakhimovich, Bludorova, Chiguryaeva *et al.* (1985), Zosimovich *et al.* (1985), Yassini (1986), Mekhtiev & Pashaly (1987), Muzylev & Golovina (1987), Steininger *et al.*, *in* Royden & Horvath (1987), Olteanu (1989), Rögl *et al.* (1991), Fedorov (1994), Markova & Mikhailesku (1994) and Jones

(1996). A comprehensive bibliography of general stratigraphic references (to 1984) is given by Rögl (1985b).

Petroleum geological aspects have been discussed by, among others, Khain *et al.* (1937), Ismailov & Idrisov (1963), Ali-Zade *et al.* (1966), Shilinski (1967), Ismailov *et al.* (1972), Buryakovsky (1974, 1993), Alikhanov (1977), Nikishin (1981), Ulmishek & Harrison (1981), Babayan (1984), Panakhi & Buare Mamadu Lamin (1987), Bagir-Zade *et al.* (1988), Akramkhodzhaev *et al.* (1989), Kerimov *et al.* (1991), Kleschev *et al.* (1992), Narimanov (1993) and Reynolds *et al.* (*in press*). Additional comments on petroleum geology are inserted as appropriate in the succeeding sections.

Maykopian (Figs 7–8)

The Maykopian takes its name from a town in the Caucasus (Likharev, 1958). The term Maykopian refers to essentially argillaceous rocks of Oligocene to Early Miocene age. The Zeivar Formation of Northern Iran and Lower Red Formation (predominantly clastics) of Central Iran (the latter locally contains age-diagnostic lepidocyclinid and nummulitid larger benthonic foraminifera) appear correlative, as does the Qom [Qum] Formation (predominantly carbonates) of Central Iran (which contains numerous age-diagnostic species of alveolinid, lepidocyclinid and miogypsinid larger benthonic foraminifera (Rahaghi, 1973)) (Stöcklin & Setudehnia, 1971, 1972). The Maykopian (in particular the Khadumian) is an important regional source rock (e.g., Veto, 1987). It also constitutes a minor reservoir in the Kobustan-Kura region of the South Caspian (Ali-Zade *et al.*, 1966).

Details of Maykopian stratigraphy have been discussed by Muratov (1960), Ali-Zade (1966), Ali-Zade & Mamedov (1970), Mamedova & Mamedova (1970), Azizbekov (1972), Lupov *et al.* (1972), Khalilov & Mamedova (1973), Bolli & Krashenninnikov (1977), Ali-Zade & Atayeva (1982), Krashenninnikov & Muzylev (1975), Krashenninnikov, Muzylev & Ptukhian (1985), Nevesskaya & Nosovsky (1985), Bugrova (1986), Koshkarly (1986, 1993), Krashenninnikov (1986), Krashenninnikov & Ptukhian (1986), Koshkarly & Baldi-Beke (1987), Gasanov & Kyazamov (1988), Nagymarosy (1992) and Koshkarly & Alekperov (1993).

Microbiostratigraphic study of the Maykopian is hindered by massive reworking, reflecting deposition in a foreland basin in front of the emerging Caucasus. Maykopian samples can contain >90% reworked (especially Eocene) microfossils.

The Maykopian has been divided into five sub-stages by various authors (see Fig. 5). In ascending stratigraphic order, these are the Khadumian, Roshenian, Caucasian, Sakaraulian and Kozakhurian. The Khadumian is dated as Early Oligocene on the evidence of planktonic foraminifera and calcareous nannofossils and can therefore be calibrated against global standard biostratigraphic zonation schemes and absolute chronostratigraphic time-scales (see below). In contrast, the Roshenian is dated as Late Oligocene and the Caucasian to Kozakhurian as Early Miocene essentially only on the evidence of benthonic foraminifera (see, for instance, Nevesskaya & Nosovsky, 1985).

The youngest sub-stages of the Maykopian appear to be absent in the Dacian Basin in the Western Black Sea region (Steininger *et al.*, *in* Royden & Horvath, 1987).

Micropalaeontology and Nannopalaeontology. The Khadumian has been dated as Early Oligocene on both planktonic foraminiferal and calcareous nannoplankton evidence. Ali-Zade




Fig. 6 Stratigraphic summary (Miocene-Holocene). Chronostratigraphy after Beard *et al.* (1982) and Lamb *et al.* (1987) (conceptual Gulf of Mexico deep-water stage nomenclature) (Wis. = Wisconsinian; San. = Sangamonian; Ill. = Illinoisian; Yar. = Yarmouthian; Kan. = Kansan; Aft. = Aftonian; Neb. = Nebraskan). Biostratigraphy after Blow (1969) (planktonic foraminifera) (prefixed N) and Martini (1971) (calcareous nannoplankton) (prefixed NN). Magnetostratigraphic polarity epochs and absolute age values are after Haq *et al.* (1988). Regional stratigraphy from this paper (upper case = chronostratigraphic, lower case = lithostratigraphic units). Regional sequence stratigraphy (transgressions) after Skálbdyna (1985). Global sequence stratigraphy after Haq *et al.* (1988) (third-order cycles TB3.2–TB3.8), and Beard *et al.* (1982) and Lamb *et al.* (1987) (fourth-order cycles Q2–Q8). Sequence boundary and maximum flooding surface ages in Ma. Climatostratigraphy (superclimathems or SCTs) after Zubakov & Borzenkova (1990). The correlation of Eastern Paratethyan sequence stratigraphy with the global coastal onlap curve of Haq *et al.* (1988) is tentative. The extent of biostratigraphic control constraining the correlation is indicated on Fig. 5.

(1966) recorded the planktonic foraminifer *Globigerina officinalis* (Middle Eocene to Oligocene, Zones P14–P22 of Blow, 1969) from the basal ‘*Planorbella* Horizon’ in Azerbaijan. Krashenninnikov (1986) recorded planktonic foraminifera indicative of Early Oligocene (P18) from a stratigraphically similar position (immediately below the ‘*Ostracod*-Horizon’) in the Kuban–Kuma Interfluvium (North Caucasus). Krashenninnikov *et al.* (1985) recorded *Globigerina tapuiensis* (Oligocene, P18–P20) associated with the larger benthonic foraminifer *Nummulites intermedius* (Early Oligocene) from the Khadumian of Armenia. Later, Krashenninnikov & Ptukhian (1986) recorded *Globigerina sellii* (Oligocene, P19/20 to ‘early’ P22) associated with *Nummulites intermedius* (also Oligocene) and the calcareous nannofossil *Helicosphaera reticulata* (Eocene to Early Oligocene, Zones NP17–NP22 of Martini, 1971) from the Khadumian of Armenia. Koshkarly (1986) recorded the

calcareous nannofossils *Reticulofenestra umbilica* (Eocene to Early Oligocene, NP16–NP22), *Chiasmolithus oamaruensis* (Eocene to Early Oligocene, NP18–NP22), *Isthmolithus recurvus* (Eocene to Early Oligocene, NP19–NP22), *Sphenolithus pseudoradians* (Eocene to Early Oligocene, NP20–NP23) and *Ericsonia subdisticha* (Eocene to Early Oligocene, NP20–NP21) from the Khadumian of Azerbaijan. Later, Koshkarly & Baldi–Beke (1987) recorded *R. umbilica*, *C. oamaruensis*, *I. recurvus* and *S. pseudoradians*, and Koshkarly & Alekperov (1993) *H. reticulata* and *E. subdisticha* from the Early Maykopian of Azerbaijan.

Palynology. Although only non-age-diagnostic palynomorphs were recorded from the Maykopian of the Middle Kura Depression by Dzhabarova (1973), recent observations suggest that some age-diagnostic dinocysts, and pollen and spores do

exist (in the Early, and Middle to Late Maykopian respectively). These will be reported in detail in a future publication. Palynological evidence indicates that the Maykopian is a regressive unit characterised by upwardly-increasing terrestrial input (upwardly-increasing pollen and spore content). Micropalaeontological and sedimentological evidence also indicates shallowing upward.

Tarkhanian to Konkian (Figs 8–10)

These stages have collectively been correlated with the Badenian of Central Paratethys (see, for instance, Steininger *et al.*, in Royden & Horvath, 1987). The Badenian is Middle Miocene (planktonic foraminiferal zones N8–?N12 (see, for instance, Papp *et al.*, 1968, Rögl *et al.*, 1978 and Papp & Schmid, 1985); calcareous nannoplankton zones NN5–NN7 (see, for instance, Rögl *et al.*, 1978, Papp & Schmid, 1985 and Meszaros, 1992)). Palynologically, it is locally characterised by mangrove elements (Nagy & Kokay, 1991).

The base of the Badenian (the Moravian sub-stage of Papp *et al.* (1978) (Lagenid Zone)) is defined at the first appearance of the planktonic foraminifer *Praeorbulina* (Zone N8) (Papp *et al.*, 1968). The first appearance of the ancestral form *Globigerinoides bisphericus*, which defines the base of Zone N8 (Blow, 1969), falls within the underlying Karpatian (Cicha *et al.*, 1967). This biostratigraphic control indicates that the base of the Badenian can be correlated with the 16.5Ma (glacio-eustatic) sea-level low-stand of Haq *et al.* (1988) (Fig. 5).

The middle part of the Badenian (the Wielician sub-stage of Papp *et al.* (1978) (Sandschaler Zone)) is characterised by marginal marine sediments (including evaporites). This

regressive sub-stage can be tentatively calibrated against planktonic foraminiferal Zones N10–N12 or calcareous nannoplankton zones NN5–NN6 (see, for instance, Rögl *et al.* (1978) and Papp & Schmid (1985)). The onset of regressive conditions can be tentatively correlated with the 15.5Ma (glacio-eustatic) sea-level low-stand of Haq *et al.* (1988) (Fig. 5). The regressive coarse clastics of the Chokrakian and Karaganian in Eastern Paratethys also appear to be associated with this event (though they could be associated with a separate tectonic event). These clastics constitute important reservoirs in the Indol Kuban and Terek Caspian Foredeeps (Ulmishek & Harrison, 1981) and in eastern Azerbaijan (Ali-Zade *et al.*, 1986). Chepalyga (1985) calibrates the Chokrakian and Karaganian against magnetostratigraphic polarity epochs 15–12, while Zubakov & Borzenkova (1990) calibrate them against polarity epochs 16–14.

The top of the Badenian (the Kosovian sub-stage of Papp *et al.* (1978) (Buliminid-Bolivinid Zone)) is defined below the last appearances of *Globorotalia mayeriisakensis* (N14) and *Globigerina druryi* (N15) (Papp *et al.*, 1978; Papp & Schmid, 1985).

Details of Tarkhanian to Konkian stratigraphy have been discussed by Andrusov (1884), Bogdanowicz (1950a–b, 1965), Shishova (1955), Gasanova (1965), Mamedova (1971), Azizbekov (1972), Lupov *et al.* (1972), Dzhabarova (1973), Cicha *et al.* (1983) and Ali-Zade *et al.* (1986). A monograph of polymorphinid foraminifera of this age from Georgia was published by Dzharelidze (1977).

In eastern Azerbaijan the Karaganian and Konkian, together with the overlying Sarmatian and Maeotian, are collectively referred to as the Diatom Suite (because of the presence of




Fig. 7 Palaeogeographic reconstruction, Early Maykopian (Early Oligocene). Solid line indicates relatively well constrained, dashed line poorly constrained shoreline. Ticks on landward side. The location of the Maykopian stratotype is indicated.

diatomaceous limestones). The individual Eastern Paratethyan stages can be recognised within the Diatom Suite on the basis on fish otoliths and benthonic microfossils (particularly molluscs and foraminifera (e.g., Azizbekov, 1972; Ali-Zade *et al.*, 1986) and diatoms (E.Z. Ateava, pers. comm., 1994)). The work of Dzhabarova (1973) suggests that palynology may also be used to recognise the various stages (see notes below).

In eastern Azerbaijan parts of the Diatom Suite are considered to have hydrocarbon source potential.

Tarkhanian (Fig. 8)

The Tarkhanian takes its name from a promontory in the Crimea (Likharev, 1958). The stratotype section yields Middle Miocene (NN5) calcareous nannoplankton (F. Rögl, pers. comm., 1994).

Micropalaeontology. Only non-age-diagnostic quasi-marine, smaller benthonic and rare planktonic foraminifera were recorded by Bogdanowicz (1950a) from the Tarkhanian of Kuban and later by Mamedova (1971) and Azizbekov (1972) from the Tarkhanian of Azerbaijan. These include *Rotalia* [*Ammonia*] ex gr. *beccarii* (smaller benthonic), which has a cosmopolitan distribution and probably ranges no older than Middle Miocene (RWJ's unpublished observations), and *Nonion* [*Florilus*] *boueanum* (smaller benthonic) and *Globigerina tarchanensis* (planktonic), both of which have also been recorded in the Badenian of Central Paratethys (Papp *et al.*, 1978; Papp & Schmid, 1985).

Chokrakian (Fig. 9)

The Chokrakian takes its name from a lake in the Crimea (Likharev, 1958). It is of Middle Miocene age on regional evidence (see above). Direct biostratigraphic evidence is lacking. The 'Vindobonian Marls' of Northern Iran appear correlative (Stöcklin & Setudehnia, 1971, 1972).

Micropalaeontology. Only non-age-diagnostic, quasi-marine, smaller benthonic foraminifera were recorded by Bogdanowicz (1950b) from the Chokrakian of the western Precaucasus and later by Mamedova (1971) and Azizbekov (1972) from the Chokrakian of Azerbaijan and Popkhadze (1983) for the Chokrakian of western Georgia. These include *Rotalia* [*Ammonia*] ex gr. *beccarii* (smaller benthonic), which has a cosmopolitan distribution and probably ranges no older than Middle Miocene (RWJ's unpublished observations), *Nonion* [*Florilus*] *boueanum* and *Miliolina* [*Quinqueloculina*] *akneriana* spp., both of which have also been recorded in the Badenian of Central Paratethys (Papp & Schmid, 1985), and *Miliolina caucasica*, *Sigmoilina tschokrakensis* and *Tschokrakella longiuscula*, all of which are endemic to Eastern Paratethys. The ostracod, *Leptocythere bardrakensis* was recorded by Popkhadze (1984) from the Chokrakian of western Georgia.

Palynology. Only non-age-diagnostic palynomorphs were recorded by Dzhabarova (1973) from the Chokrakian of the Middle Kura Depression. Pollen spectra are characterised by relatively high incidences of herb and shrub taxa including *Chenopodiaceae* and *Ephedra*. The presence of




Fig. 8 Palaeogeographic reconstruction. Late Maykopian to Tarkhanian (Late Oligocene to early Middle Miocene). Key as for Fig. 7. The locations of the Maykopian and Tarkhanian stratotypes are indicated.




Fig. 9 Palaeogeographic reconstruction, Chokrakian to Karaganian (Middle Miocene). Key as for Fig. 7. The location of the Karaganian stratotype is indicated.

Chenopodiaceae probably indicates the local development of salt-marshes.

Karaganian (Fig. 9)

The Karaganian takes its name from a locality on the Mangyshlak Peninsula in Kazakhstan (Likharev, 1958). It is Middle Miocene on regional evidence (see above). Direct biostratigraphic evidence is lacking. The *Spaniodontella* Beds of Northern Iran appear correlative (Stöcklin & Setudehnia, 1971, 1972).

Micropalaeontology. Only non-age-diagnostic, quasi-marine, smaller benthonic foraminifera were recorded by Mamedova (1971) and Azizbekov (1972) from the Karaganian of Azerbaijan. These include *Nonion bogdanowiczi*. The fish otoliths *Rhombus corius* and *R. corius binagadinica* are regarded as index-species for the Karaganian in Azerbaijan (E.Z. Ateava, pers. comm., 1994).

Palynology. Only non-age-diagnostic palynomorphs were recorded by Dzhabarova (1973) from the Karaganian of the Middle Kura Depression. Pollen spectra are characterised by relatively high incidences of tree taxa, which indicates a forested hinterland. The predominance of *Betula* (birch) indicates a climatic regime similar to that of the present-day taiga or forest-tundra.

Konkian (Fig. 10)

The Konkian takes its name from a river in the Ukraine (a tributary of the Dniepr) (Likharev, 1958). It is of Middle

Miocene age on regional evidence (see above). Direct biostratigraphic evidence is lacking. The *Pholas* Beds of Northern Iran appear correlative (Stöcklin & Setudehnia, 1971, 1972).

Micropalaeontology. Only non-age-diagnostic, quasi-marine, smaller benthonic foraminifera were recorded by Bogdanowicz (1965) from the Konkian of the western Precaucasus and by Mamedova (1971) and Azizbekov (1972) from the Konkian of Azerbaijan. These include *Rotalia* [*Ammonia*] ex gr. *beccarii* (smaller benthonic), which has a cosmopolitan distribution and probably ranges no older than Middle Miocene (RWJ's unpublished observations), *Articulina gibbosa* and *Miliolina* [*Quinqueloculina*] *haidingerii*, both of which have also been recorded in the Badenian of Central Paratethys (Papp & Schmid, 1985), and *Articulina elongata konkensis*, *Bulimina konkensis* and *Elphidium nachischevanicus*, all of which are endemic to Eastern Paratethys. *Bulimina konkensis* and *Elphidium kudakoense*, together with the fish otolith *Trigla konkensis*, are regarded as index-species for the Konkian in Azerbaijan (Podobina *et al.*, 1956; Mamedova, 1971).

Shishova (1955) and Gasanova (1965) recorded the following diatoms from the Konkian of Eastern Azerbaijan: *Actinocyclus ehrenbergi*, *A. rafsii*, *Asterolampra marylandica*, *Cocconeis placentula lineta*, *C. scutellum*, *Coscinodiscus radiatus*, *C. oculus* and *Melosira sulcata*. *Coscinodiscus radiatus* was considered particularly typical.

Palynology. Only non-age-diagnostic palynomorphs were recorded by Dzhabarova (1973) from the Konkian of the Middle Kura Depression. Pollen spectra are characterised by relatively high incidences of tree taxa, which indicates a forested




Fig. 10 Palaeogeographic reconstruction. Konkian to Maeotian (late Middle to early Late Miocene). Key as for Fig. 7. The location of the Konkian stratotype is indicated.

hinterland. The predominance of Taxodiaceae (cypresses and swamp-cypresses) indicates a warm-temperate (possibly even subtropical) climatic regime.

Sarmatian (Fig. 10)

The Sarmatian of Eastern Paratethys is probably equivalent to the stratotypical Sarmatian of Central Paratethys (late Middle to early Late Miocene (calcareous nannoplankton zones NN7–NN9) (Meszaros, 1992), planktonic foraminiferal zones N13?–N15), but may also be equivalent to the lower part of the Pannonian (Slavonian) of that area (Late Miocene) (see, for instance, Papp *et al.*, 1974, 1985). Chepalyga (1985) calibrates the Sarmatian of Eastern Paratethys against magnetostratigraphic polarity epochs 10–7, while Zubakov & Borzenkova (1990) calibrate it against epochs 14–9, and Pevzner & Vangengeim (1993) calibrate it against polarity epochs 10–7. The regressive events which characterise the Sarmatian suggest that (within the limits of biostratigraphic control) its base can be correlated with the 10.5Ma (glacio-eustatic) sea-level low-stand of Haq *et al.* (1988).

The Sarmatian of Eastern Paratethys is characterised by areally restricted regressive marginal marine sediments including coarse clastics, and, in the Nakchichevan Depression, evaporites (though transgressive black shales (with source potential) also occur locally). The 'Sarmatian' of Northern Iran and part of the Upper Red Formation of Central Iran, also characterised by clastics and evaporites, appear correlative (Stöcklin & Setudehnia, 1971, 1972). The Sarmatian of Eastern Paratethys has been divided into three sub-stages, which are, from oldest to youngest, Volkhyanian, Bessarabian and

Chersonian. The 'mid' Sarmatian (Bessarabian) represents the culmination of a major regressive phase that began in late Konkian or 'early' Sarmatian (Volkhyanian) times (see, for instance, Chepalyga, 1985), and resulted in the first isolation of the South Caspian Basin.

Details of the Sarmatian stratigraphy of Eastern Paratethys have been discussed by Gasanova (1965), Maisuradze (1971), Mamedova (1971, 1987), Azizbekov (1972), Lupov *et al.* (1972), Dzhabarova (1973), Ali-Zade & Aleskerov (1974), Azizbekova (1974), Paramonova *et al.* (1979) and Pevzner & Vangengeim (1993).

Micropalaeontology. Only non-age-diagnostic (and largely endemic), quasi-marine, smaller benthonic foraminifera and ostracods were recorded from the Sarmatian by Podobina *et al.* (1956), Mamedova (1971, 1987), Voroshilova (1971) and Azizbekov (1972) from Azerbaijan, by Azizbekova (1974) from the Nakchichevan Depression, and by Paramonova *et al.* (1979) from various sites in the Ponto-Caspian region. These include rare cosmopolitan species such as *Streblus* [*Ammonia*] *beccarii* and *Elphidium macellum* (foraminifera), which probably range no older than Middle Miocene (RWJ's unpublished observations), *Elphidium reginum* (foraminifer) (lower part only), which is also found in the Sarmatian of Central Paratethys (Steininger *et al.*, 1976) and (?reworked) in the Pliocene of the Black Sea (Gheorghian, in Ross *et al.*, 1978), *Nonion* div. spp., *Porosonion* div. spp. and *Quinqueloculina consobrina* (foraminifera) and *Cyprideis littoralis*, *Cythere multistriata*, *Leptocythere stabilis*, *Loxoconcha eichwaldi* and *Xestoleberis lutrae* (ostracods). Podobina *et al.* (1956), Mamedova (1971) and Voroshilova (1971) have indicated that the three subdivisions of

the Sarmatian in Azerbaijan can be recognised on the basis of foraminifera and ostracods.

The alga *Ovulites sarmaticus* and the fish otoliths *Gadidarum minusculus* and *Gobius sarmaticus* are regarded as index-species for the Sarmatian in Azerbaijan (Ateava, personal communication, 1994). The Sarmatian part of the 'Diatom Suite' in Azerbaijan also contains 23 species of diatoms, chiefly *Coscinodiscus*, *Licmophora* and *Navicula* (Gasanova, 1965).

Palynology. Only non-age-diagnostic palynomorphs were recorded from the Sarmatian of the Middle Kura Depression (Azizbekov, 1972; Dzhabarova, 1973). Pollen spectra from the middle (Bessarabian) sub-stage are characterised locally by (as in the 'Cryptomactra-Horizon') relatively high incidences of Pinaceae (pine) pollen and locally (higher in the section) by relatively high incidences of broad-leaved tree pollen (*Alnus* (alder), *Betula* (birch), *Carya* (hickory), *Carpinus* (hornbeam), *Corylus* (hazel), *Juglans* (walnut)) and angiosperm (flowering plant) pollen (Azizbekov, *op. cit.*). Pollen spectra from the upper (Chersonian) sub-stage are also characterised locally by relatively high incidences of broad-leaved tree pollen (*Fagus* (beech), *Quercus* (oak), *Taxodium* (swamp-cypress), *Tilia* (lime) and *Ulmus* (elm)), though locally Polypodiaceae predominate (Azizbekov, *op. cit.*).

Maeotian (Fig. 10)

The Maeotian of Eastern Paratethys is equivalent to the Pannonian (Late Miocene) of Central Paratethys (though possibly only the upper part thereof (Serbian) (see, for instance, Papp *et al.* (1974, 1985) and Pevzner & Vangengeim (1985b)). Rögl (1985a) calibrates it against calcareous nannoplankton Zone NN10. Chepalyga (1985) calibrates it against magnetostratigraphic polarity epochs 6–5, while Zubakov & Borzenkova (1990) calibrate it against polarity epochs 10–7. Krakhmalnaya *et al.* (1993) have recently described a succession at Novaya Emetovka on the Black Sea coast with a good Maeotian mammal fauna which they calibrate against polarity epochs 6 and 5. The Maeotian is represented by a transgressive-regressive cycle. It is locally characterised by black shales (with hydrocarbon source potential). The Maeotian sea was probably characterised by reduced salinity. Similar environmental conditions evidently obtained in the Pannonian of Central Paratethys, where benthonic foraminiferal assemblages are of quasi-marine (brackish water) aspect (*Ammobaculites*, *Ammomarginulina*, *Miliammina*, *Trochammina*) (Papp *et al.*, 1985).

Details of Maeotian stratigraphy have been discussed by Shishova (1955), Gasanova (1965), Bogdanowicz (1967, 1969, 1974), Mamedova (1971), Voroshilova (1971), Azizbekov (1972), Lupov *et al.* (1972), Popkhadze (1977), Paramonova *et al.* (1979), Ananova *et al.* (1985), Ali-Zade *et al.* (1986), Rasulov (1986), Maisuradze (1988), Naidina (1988) and Ateava (personal communication, 1994).

Micropalaeontology and Nannopalaeontology. Only non-age-diagnostic, quasi-marine, smaller benthonic foraminifera were recorded by Bogdanowicz (1967, 1969) from the Maeotian of the Kuban and Western Precaucasus and by Paramonova *et al.* (1979) from the Maeotian of various sites in the Ponto-Caspian. These include *Elphidium macellum*, which has a cosmopolitan distribution and probably ranges no older than Middle Miocene (RWJ's unpublished observations) and *Nonion* div. spp.

Some stratigraphically and/or palaeoenvironmentally

significant ostracods were recorded by Azizbekov (1972) from Azerbaijan and by Popkhadze *et al.* (1980) from Abkhazia. These include *Leptocythere biplicata*, *L. meotica*, *Loxoconcha meotica*, *L. tamarindus* and *L. viridis* (quasi-marine). *Leptocythere meotica* and *Loxoconcha meotica*, together with the foraminifera *Quinqueloculina sulacensis* (lower part) and *Q. ludwigi* (upper part) and the fish otoliths *Clupea gidjakensis* and *Percidarium sigmalinoides* are regarded as index-species for the Maeotian in Azerbaijan (Podobina *et al.*, 1956; Mamedova, 1971).

Rich diatom floras including *Coscinodiscus gigas*, *C. oclisirides*, *Grammatophora azens*, *Cocconeis heteroidea*, *Melosira architecturalis* and *M. sulcata* were recorded by Shishova (1955) from the Maeotian part of the 'Diatom Suite' of the Apsheron Peninsula in Azerbaijan. *Actinocyclus ehrenbergii*, *Cymatosira sovitchenkoi* and *Rhaphoneis maeotica* were recorded by Gasanova (1965). The stenohaline (normal marine) forms *Asterolampha marylanica*, *Coscinodiscus asteromphalus* and *C. lewisianus sensu lato*, the euryhaline (quasi-marine) forms *Actinocyclus ehrenbergii* and *Rhapalodia musculus*, and the freshwater forms *Amphora ovalis* and *Diatoma vulgare* were recorded by Rasulov (1986). *Coscinodiscus lewisianus sensu stricto* is a Middle Miocene species.

A range of diatoms, calcareous nannofossils and ostracods were recorded by Ananova *et al.* (1985) from the Maeotian of the Black Sea. These include *Actinocyclus ehrenbergii*, *Rhaphoneis maeotica* and *Thalassiosira maeotica* (diatoms), *Braarudosphaera* spp. (calcareous nannofossil) and *Cyprideis torosa* and *Leptocythere* spp. (ostracods).

Palynology. Only non-age-diagnostic palynomorphs have been recorded from the Maeotian, and only the acritarch *Micrhystridium* sp. was recorded by Ananova *et al.* (1985) from the Maeotian of the Black Sea. Pollen spectra from the Late Maeotian of the Task-Sunzhenskii Region are characterised by relatively high incidences of Asteraceae and Polygonaceae (herbs), *Ephedra* (shrubs) and Gramineae (grasses) (Naidina, 1988). This indicates an open, sparsely forested hinterland and an arid climatic regime similar to that of the present-day steppe or semi-desert. The locally relatively high incidences of Gleicheniaceae (ferns) and Lycopodiaceae (club-mosses) indicate local development of conditions similar to those of the present-day tundra, forest-tundra or mountain belt. The presence of Chenopodiaceae probably indicates local development of salt-marshes.

Pontian

The Pontian takes its name from the ancient name for the Black Sea. It is essentially regressive (though it also includes an overstepping transgressive unit at the base), and is characterised regionally by marginal marine coarse clastics and shallow marine carbonates and locally (Babajan Formation, Bosporian Sub-Stage) by evaporites. Chepalyga (1985) calibrates the Pontian against magnetostratigraphic polarity epoch 4 (Gilbert), while Zubakov & Borzenkova (1990) calibrate it against polarity epochs 6–5. We correlate the Pontian evaporites against those of the stratotypical Messinian, which can be calibrated against magnetostratigraphic polarity epoch 5 (Zubakov & Borzenkova, 1990).

Details of the Pontian stratigraphy of Eastern Paratethys have been discussed by, among others, Sveier (1949), Mandelstam *et al.* (1962), Sheydayeva-Kuliyeva (1966), Agalarova (1967), Vekilov *et al.* (1969), Ramishvili (1969), Rozyeva (1971),

Azizbekov (1972), Lupov *et al.* (1972), Chelidze (1973), Karmishina (1975), Vekua (1975), Krstic (1976), Shchekina (1979), Imnadze & Karmishina (1980), Ali-Zade *et al.* (1986), Sirenko & Turlo (1986) and Yagmurulu & Helvacı (1994).

Micropalaeontology. Only the non-age-diagnostic, quasi-marine, smaller benthonic foraminifer *Elphidium stellatum* was recorded by Imnadze & Karmishina (1980) from the Late Pontian (Bosporian sub-stage) of the Black Sea.

Stratigraphically and/or palaeoenvironmentally significant ostracods recorded by Sheydayeva-Kuliyeva (1966) and Azizbekov (1972) from the Pontian of Azerbaijan (Western Caspian), by Karmishina (1975) from the Northern Precaspian and South-Eastern Kalmyk (Northern Caspian) and Prechernomore (Northern Black Sea), and by Vekua (1975) from Abkhazia (north-eastern Black Sea) include *Bakunella dorsoacuta*, *Caspiolla acronasuta*, *Pontoniella acuminata* and *P. loczyi*. *Bakunella dorsoacuta* and *Caspiolla acronasuta* are quasi-marine species (Yassini, 1986). Species of *Leptocythere*, *Loxoconcha* and *Xestolebris* also occur. It is possible to recognise three ostracod zones in the Pontian of Eastern Azerbaijan (Sheydayeva-Kuliyeva, 1966). It is also possible to recognise three corresponding mollusc zones.

The lower part of the Continental (Cheleken) Series of Northern Iran and the Maragheh Bone Beds of Central Iran can be correlated with the Pontian on vertebrate palaeontological, limited malacological and ostracod evidence (Faridi, 1964; Stöcklin & Setudehnia, 1971, 1972).

Palynology. Only non-age-diagnostic palynomorphs have been recorded from the Pontian. Pollen spectra from the Pontian of Georgia are characterised by relatively high incidences of tropical elements such as *Nypa* (palm) (Ramishvili, 1969). Those

from the Late Pontian (Bosporian sub-stage) of the Ukraine are characterised initially by thermophilic (warm-temperate) and hydrophilic (moisture-loving) elements such as Taxodiaceae (cypresses and swamp-cypresses) and later by arid steppe and semi-desert elements (Shchekina, 1979; Sirenko & Turlo, 1986).

Kimmerian (Fig. 11)

The Kimmerian takes its name from an ancient tribe who lived on the shores of the Black Sea (Likharev, 1958). The Dacian (a stage name sometimes used in the Black Sea region) and the 'Eoakchagyl' (a stage name used in the Caspian Sea region by Zubakov & Borzenkova (1990)) appear synonymous. Semenenko (1979), Pevzner & Vangengeim (1985), Skalbldyna (1985) and Zubakov & Borzenkova (1990) calibrate the Kimmerian against magnetostratigraphic polarity epochs 5–4, while Chepalyga (1985) and Chepalyga *et al.* (1985) calibrate it against polarity epoch 4 (Gilbert). Essentially on the basis of magnetostratigraphic evidence (including the calibration of the underlying Pontian against polarity epoch 5 (see above)), we have tentatively calibrated the major unconformity at the base of the Kimmerian against the 5.5Ma glacio-eustatic sea-level low-stand of Haq *et al.* (1988) and apparently coincident uplift around and subsidence within the Caspian (leading to a massive sea-level fall (of the order of 1000m) within the Caspian and the severance of the connection between the Caspian and the Black Sea). Additional unconformities in the Caspian succession cannot be confidently calibrated against any of the third-order glacio-eustatic sea-level low-stands on the Haq *et al.* chart, and may be associated with higher frequency glacio-eustatic low-stands or local tectonic events. Due partly to tectono-eustatic effects (see above), and partly to climatic effects




Fig. 11 Palaeogeographic reconstruction, Dacian/Kimmerian (latest Miocene to Pliocene). Key as for Fig. 7.

(an excess of evaporation over precipitation) and a change in the drainage system (with rivers formerly flowing into Paratethys captured by rejuvenated rivers flowing into the Mediterranean, where base-level had fallen considerably during the Messinian salinity crisis (see, for instance, Hsu (1983)), the level and the areal extent of the Caspian 'athalassic lake' were drastically reduced during the Kimmerian. Palaeontological evidence points to a stratigraphically upward reduction in salinity. The basal Kimmerian contains a quasi-marine ostracod fauna (characterised by *Cyprideis littoralis/torosa*) that appears correlative with that of the Messinian 'Lago Mare' facies in the Mediterranean.

In Azerbaijan, the Kimmerian is characterised by several thousand metres (thicknesses of around 4 km are typical around the Apsheron Peninsula) of principally Palaeo-Volga-derived, regressive marginal and non-marine coarse clastics comprising the main ('Productivnaya Tolsha' = 'Productive Series') reservoirs in several billion-barrel oil-fields (Khain *et al.*, 1937; Ismailov and Idrisov, 1963; Ismailov *et al.*, 1972; Nikishin, 1981; Babayan, 1984; Kerimov *et al.*, 1991; Narimanov, 1993). The sedimentology of this succession is discussed in detail by Reynolds *et al.* (in press). Constituent lithostratigraphic units include, in ascending stratigraphic order, the Kalin or Kala, Podkirmakina (Pre-Kirmaky Sand (PK)), Kirmakina (Kirmaky Sand (KS)), Nadkirmakina (Post-Kirmaky Sand and Post-Kirmaky Clay (NKP and NKG)), Pereriva, Balakhany, Sabunchi and Surakhany Suites.

Details of Kimmerian stratigraphy have been discussed by, among others, Ali-Zade (1946), Khalilov (1946), Sveier (1949), Agalarova (1956), Mandelstam *et al.* (1962), Rozyeva (1971), Azizbekov (1972), Lupov *et al.* (1972), Ali-Zade (1974), Karmishina (1975), Vekua (1975), Ananova *et al.* (1985), Sirenko & Turlo (1986) and Yassini (1986).

Micropalaeontology Only non-age-diagnostic, quasi-marine, smaller benthonic foraminifera were recorded by Karmishina (1975) from the Kimmerian of Prechernomore (Northern Black Sea). These include *Ammonia beccarii* and *Elphidium incertum*.

Stratigraphically and/or palaeoenvironmentally significant ostracods recorded by Karmishina (1975) and Vekua (1975) from the Kimmerian of Prechernomore (Northern Black Sea) and Abkhazia (North-eastern Black Sea) respectively include *Bakunella dorsoacuata*, *Caspiocypris labiata*, *Caspiolla acronasuta*, *Cyprideis littoralis/torosa*, *Leptocythere bosqueti*, *Mediocythereis apatoica* and *Pontoniella acuminata*. *Bakunella dorsoacuata*, *Caspiolla acronasuta* and *Cyprideis torosa* are quasi-marine species (Yassini, 1986). *Caspiocypris labiata* is fresh-water. Ostracods recorded by Karmishina (1975) from the Kimmerian of the Northern Precaspian and South-Eastern Kalmyk (Northern Caspian) include *Candona angulata*, *C. neglecta*, *C. rostrata*, *Cyclopypris laevis*, *Cypria arma*, *Eucypris naidinae*, *Ilyocypris bradyi*, *I. gibba*, *I. serpulosa* and *Zonocypris membranacea*. These are all fresh-water forms.

Microbiostratigraphic study of the Kimmerian 'Productive Series' in Azerbaijan is hindered by massive reworking (Khalilov, 1946; Agalarova, 1956). However, the general pattern of reworking indicates unroofing, which in itself is of some stratigraphic value (with pulses of reworking related to sea-level low-stands). Moreover, the potential exists for biostratigraphic subdivision on the basis of ostracods (see, for instance, Khalilov, 1946; Agalarova, 1956; Azizbekov, 1972). The Kala Suite contains *Cythere* [*Leptocythere*] *camellii* and *Paracypris liventalina* (quasi-marine), the Kirmakina Suite *Cythere* [*Leptocythere*] *cellula*, *C. [Cyprideis] littoralis/torosa*, *C. olivina*,

C. [Leptocythere] praebacuana, *Hemicythere pontica* and *Loxoconcha djabaroffi* (quasi-marine), the Pereriva Suite *Cyprideis littoralis/torosa* (quasi-marine), the Balakhany Suite *Cythere* [*Leptocythere*] *praebacuana*, *Loxoconcha alata*, *L. eichwaldi* and *Xestoleberis lutrae* (quasi-marine), the Sabunchi Suite no *in situ* ostracods, and the Surakhany Suite *Eucypris* (fresh-water) and *Leptocythere* and *Loxoconcha* (quasi-marine).

The Krasnotsvetnaya ('Red-Coloured') Series of Turkmenia can be correlated with the Kimmerian on ostracod evidence (Agalarova, 1956; Lupov *et al.*, 1972). The lowermost part contains *Bakunella*, *Caspiolla*, *Pontoniella* and *Xestoleberis* and is of Pontian aspect, while the uppermost part contains *Leptocythere* and *Limnocythere* and is of Akchagylia aspect.

The upper part of the Continental (Cheleken) Series of Northern Iran can be correlated with the Kimmerian on vertebrate palaeontological, limited malacological and ostracod evidence (Faridi, 1964; Stöcklin & Setudehnia, 1971, 1972). The Lacustrine Fish Beds of Central Iran, locally characterised by volcanic ash bands, can also be tentatively correlated with the Kimmerian (Stöcklin & Setudehnia, *op. cit.*).

Nannopalaeontology. Stratigraphically significant calcareous nannofossils recorded by Zubakov & Borzenkova (1990) from the Kimmerian include *Discoaster quinqueramus* (NN11) (?reworked), *Ceratolithus acutus* ('late' NN12) and *C. rugosus* (NN13-?NN19).

Palynology. Only non-age-diagnostic palynomorphs have been recorded from the Kimmerian. However, at least theoretically, climatostratigraphy ought to have some utility in the stratigraphic subdivision of the 'Productive Series' (with a number of superclimathems discernable on the basis of different pollen spectra). Indeed, the results of preliminary analyses have shown pollen spectra from 'Thermo'-SCT27 in the Ukraine to be characterised by broad-leaved and subtropical forest elements (*Castanea* (chestnut), *Rhus* (rue) and *Taxodiaceae* (cypress and swamp-cypress) (Sirenko & Turlo, 1986), and those from 'Cryo'-SCT26 and 'Cryo'-SCT24 to be characterised by steppe elements (Ananova *et al.*, 1985; Zubakov & Borzenkova, 1990). 'Thermo'-SCT23 has been shown to be characterised by the quasi-marine acritarch *Sigmallina* (and the quasi-marine diatom *Actinocyclus ehrenbergi*) (Ananova *et al.*, Zubakov & Borzenkova, *op. cit.*).

Akchagylia to Khvalynian (Caspian Sea) (Fig. 12)

The Akchagylia to Khvalynian of the Caspian Sea appear equivalent to the Kuyalnikian to Neoeuxinian of the Black Sea (Late Pliocene to Holocene). Details of Akchagylia to Khvalynian stratigraphy have been discussed by, among others, Livalent (1929), Sveier (1949), Mandelstam *et al.* (1962), Yakhimovich *et al.* (1965, 1984), Dzhabarova (1966), Kuliyeva (1968), Dmitriyev *et al.* (1969), Rayevskiy (1969), Sultanov & Sheydayeva-Kuliveya (1969), Aliyev & Aliyeva (1970), Kopp (1970), Rozyeva (1971), Ushakova & Ushko (1971), Zubakov & Kochegura (1971), Azizbekov (1972), Lupov *et al.* (1972), Ananova (1974), Karmishina (1975), Kaplin (1977), Trubikhin (1977), Fedkovich (1978), Veliyev (1980), Abramova (1982, 1985), Semenenko & Lulieva (1982), Mamedov & Rabotina (1984a), Aliyulla *et al.* (1985), Ivanova (1985), Mamedov & Aleskerov (1985, 1986), Sultanov (1985), Yassini (1986), Aleskerov *et al.* (1987), Bludorova *et al.* (1987), Zhidovinov *et al.* (1987), Naidina (1988), Ali-Zade & Aliyeva (1989), Yanko (1990a-b, 1991), Trubikhin *et al.* (1991) and Mamedova (1993).

Akchagylian (Fig. 12)

The Akchagylian takes its name from a locality near Krasnovodsk on the southern shores of the Gulf of Karabogaz in Western Turkmenia (Likharev, 1958). It appears equivalent to the Kuyalnikian of the Black Sea. Zubakov & Borzenkova (1990) calibrate it against magnetostratigraphic polarity epochs 4 (Gilbert) or 3 (Gauss) to 2 (Matuyama). The Akchagylian is transgressive and is represented by marine shales (constituting an important regional seal in the South Caspian). The transgressions within the Akchagylian temporarily re-established marine connections to the world's oceans. Calcareous nannoplankton data (see below) indicates that they may correspond to the 3.4Ma, 2.7Ma and 2.0Ma maximum flooding surfaces of Haq *et al.* (1988).

Details of Akchagylian stratigraphy have been discussed by, among others, Ali-Zade (1936), Yakhimovich *et al.* (1965, 1983), Dzhabarova (1966), Ali-Zade (1969), Dmitriyev *et al.* (1969), Sultanov & Sheydayeva-Kuliyeva (1969), Aliyev & Aliyeva (1970), Ushakova & Ushko (1971), Zubakov & Kochegura (1971), Azizbekov (1972), Lupov *et al.* (1972), Ananova (1974), Karmishina (1975), Trubikhin (1977), Fedkovich (1978), Sultanov (1979), Semenenko & Lulieva (1982), Mamedov & Rabotina (1984a), Bludorova *et al.* (1987), Zhidovinov *et al.* (1987), Naidina (1988) and Benyamovskoy & Naidina (1990).

Micropalaeontology. Only non-age-diagnostic smaller benthonic foraminifera were recorded by Yakhimovich *et al.* (1965), Sultanov & Sheydayeva-Kuliyeva (1969) and Karmishina (1975) from the Akchagylian of the Volga-Urals Region, Azerbaijan, and the Northern Precaspian and South-Eastern Kalmyk (Northern Caspian) respectively. These

include *Ammonia beccarii*, *Bolivina* ex gr. *advena*, *B. aksaica*, *B. floridana*, *Buccella* sp., *Cassidulina crassa*, *C. oblonga*, *Cassidulinita prima*, *Cibicides lobatulus*, *Discorbis arculus*, *D. multicameratus*, *D. orbicularis*, *D. pliocenicus*, *Elphidium incertum*, *E. kudakoense*, *E. macellum*, *E. subarcticum*, *Nonion aktschagylicus* and *Quinqueloculina* spp. *Bolivina*, *Cibicides* and *Discorbis* are near-normal marine genera, while the remainder are quasi-marine.

Stratigraphically and/or palaeoenvironmentally significant ostracods recorded by Sultanov & Sheydayeva-Kuliyeva (1969) and Karmishina (1975) from the Akchagylian of Azerbaijan, and the Northern Precaspian and South-Eastern Kalmyk (Northern Caspian) respectively include *Aglaioypris chuzchievae*, *Candona convexa*, *Ilyocypris gibba* and *Leptocythere verrucosa*. *Aglaioypris*, *Candona* and *Ilyocypris* are fresh-water forms, while *Leptocythere* is quasi-marine.

In Azerbaijan, the Akchagylian (Akchagyl Suite) contains a diverse marine ostracod and molluscan fauna (Livental, 1929; Ali-Zade, 1954; Sultanov & Sheydayeva-Kuliyeva, 1969). On the Apsheron Peninsula, it can be divided into three units on the basis of ostracods. Freshwater forms (*Candona*, *Eucypris*, *Ilyocypris*, *Limnocythere*) characterise the lower unit, marine and quasi-marine forms the middle and upper units.

Stratigraphically and/or palaeoenvironmentally significant diatoms recorded by Ushakova & Ushko (1971) from the Akchagylian to Apsheronian of the Krasnovodsk Peninsula in Western Turkmenia include *Actinocyclus ehrenbergi* ssp. *ralfsii*, *Campylodiscus noricus*, *C. noricus* ssp. *hibernica*, *Cocconeis scutellum*, *Coscinodiscus argus*, *Cymbatopleura elliptica*, *Diploneis bombus*, *D. domblittensis*, *D. fusca* ssp. *pervasta*, *D. rombica*, *Epithemia turgida*, *Grammatophora oceanica*, *Melosira*



Fig. 12 Palaeogeographic reconstruction, Akchagylian/Kuyalnikian (Late Pliocene). Key as for Fig. 7. The location of the Akchagylian stratotype is indicated.

arenaria, *M. scabrosa*, *M. sulcata*, *Navicula lyra* ssp. *elliptica*, *Nitzschia cocconeiformis*, *Rhopalodia parallella*, *Surirella striatula*, *Terpsinoe musica* and *Triceratium* sp. *Campylodiscus noricus*, *C. noricus* ssp. *hibernica*, *Diploneis domblittensis*, *D. rombica*, *Melosira arenaria*, *M. scabrosa*, *M. sulcata* and *Rhopalodia parallella* are exclusively fresh-water forms, while the remainder are quasi-marine.

The Akchagyl Beds of Northern Iran can be correlated with the stratotypical Akchagyl on ostracod evidence (Faridi, 1964; Stöcklin & Setudehnia, 1971, 1972). Various volcanics in Central Iran can also be tentatively correlated with the Akchagyl (Stöcklin & Setudehnia, *op. cit.*).

Nannopalaeontology. Stratigraphically significant calcareous nannofossils recorded by Semenenko & Lulieva (1982) from the Akchagyl include *Discoaster brouweri* (NN8?–NN18) (no younger than 1.89Ma), *D. pentaradiatus* (NN9?–NN17) (no younger than 2.33–2.43Ma) and *Reticulofenestra pseudumbilica* (NN7?–NN15). These are similar to nannoflora found in the Kuyalnikian of the Black Sea region by Semenenko & Pevzner (1979), and suggest a link between the two basins at this time.

Palynology. Only non-age-diagnostic palynomorphs have been recorded from the Akchagyl. Pollen spectra from 'Cryo'-SCT14 (and 'Thermo'-SCT15) are characterised by high incidences of *Abies* (fir), indicating a forested hinterland similar to that of the present-day taiga, and a cool, wet climate (Bludorova *et al.*, 1987). Those from 'Thermo'-SCT13 are characterised by *Abies* (fir), *Acer* (maple), *Carpinus* (hornbeam) and *Ulmus* (elm), indicating a warmer climate (Bludorova *et al.*, *op. cit.*). Pollen spectra from 'Cryo'-SCT14 in the Volga-Urals Region are characterised by relatively high incidences of Lycopodiaceae (club-mosses), again indicating a cool, wet climate (Yakhimovich *et al.*, 1965, 1983, 1984). Pollen spectra from 'Cryo'-SCTs 12 and 10 are characterised by alternations of *Alnus* (alder), *Betula* (birch) and *Pinus* (pine), indicating a cool, dry climate, and *Tsuga* (hemlock), indicating a warmer, wetter climate (Ananova, 1974; Bludorova *et al.*, 1987). Those from 'Thermo'-SCT11 in the Volga Basin are characterised by relatively high incidences of *Tsuga* (hemlock) (Zhidovinov *et al.*, 1987). Pollen spectra from the undifferentiated Akchagyl of the Tersko-Sunzhenskii Region are characterised by high incidences of broad-leaved tree taxa such as *Fagus* (beech) and *Quercus* (oak), indicating a forested hinterland similar to that of the present-day taiga (Broad-Leaved Forest Zone), and a warm-temperate climatic regime (Naidina, 1988).

Apsheeronian

The Apsheeronian takes its name from the Apsheeron Peninsula in Eastern Azerbaijan (Likharev, 1958). It appears equivalent to the Gurian of the Black Sea (Pleistocene). Zubakov & Borzenkova (1990) calibrate it against magnetostratigraphic polarity epoch 2 (Matuyama). The Apsheeronian is essentially regressive in character. Marine connections between the Caspian and the Black Sea were probably only intermittently developed.

Details of Apsheeronian stratigraphy have been discussed by Livental (1929), Sultanov (1964), Kuliyeva (1968), Ushakova & Ushko (1971), Azizbekov (1972), Lupov *et al.* (1972), Ali-Zade (1973), Ivanova (1985), Zhidovinov *et al.* (1987) and Mamedova (1988).

Micropalaeontology. Stratigraphically and/or palaeoenviron-

mentally significant ostracods recorded by Kuliyeva (1968) and Mamedova (1984, 1988) from the Apsheeronian of the Baku Archipelago in Azerbaijan include *Bakunella dorsoacuata*, *Candona albicans*, *Caspiocypris lyrata*, *C. rotulata*, *C. sinistrolyrata*, *Caspiolla acronasuta*, *C. gracilis*, *Eucypris membranae*, *Leptocythere andrusovi*, *L. caspia*, *L. fragilis*, *L. multituberculata*, *L. propinqua*, *L. quinquetuberculata*, *L. turquianica*, *L. verrucosa*, *Loxococoncha eichwaldi*, *L. gibboida*, *L. impressa* and *Trachyleberis azerbaijdzhanica*. *Bakunella*, *Caspiolla*, *Leptocythere*, *Loxococoncha* and *Trachyleberis* are quasi-marine forms, while *Candona*, *Caspiocypris* and *Eucypris* are fresh-water (see, for instance, Yassini, 1986). At least six assemblage zones can be recognised on the basis of ostracods (D.N. Mamedova, pers. comm., 1994). These will be discussed in a forthcoming paper (Mamedova, in press) (see also Mamedova, 1988).

Stratigraphically and/or palaeoenvironmentally significant diatoms recorded by Ushakova & Ushko (1971) from the Akchagyl to Apsheeronian of the Krasnovodsk Peninsula in Western Turkmenia are listed under 'Akchagyl' above. They include exclusively fresh-water, fresh-water to brackish, and brackish to near-normal marine forms.

The Apsheeron Beds of Northern Iran can be correlated with the stratotypical Apsheeronian on ostracod evidence (Faridi, 1964; Stöcklin & Setudehnia, 1971, 1972).

Palynology. Only non-age-diagnostic palynomorphs have been recorded from the Apsheeronian. Pollen spectra described by Ivanova (1985) from the Apsheeronian of Western Turkmenia and by Zhidovinov *et al.* (1987) from the Apsheeronian of the Volga Basin are characterised by relatively high incidences of non-tree taxa. This indicates an open, sparsely forested hinterland similar to the present-day steppe or forest-steppe, and a cool-temperate climatic regime.

Bakunian

The Bakunian takes its name from the city of Baku in Eastern Azerbaijan (Likharev, 1958). It appears equivalent to the Chaudian of the Black Sea (Pleistocene). Zubakov & Borzenkova (1990) calibrate it against magnetostratigraphic polarity epochs 2 (Matuyama) to 1 (Brunhes).

Details of the Bakunian stratotype have recently been discussed by Mamedova (1984, 1985, 1993, and in press) and Aliyulla *et al.* (1985). Stratigraphically and/or palaeoenvironmentally significant ostracods recorded by Aliyulla *et al.* (*op. cit.*) include *Bakunella dorsoacuata*, *Caspiocypris filona*, *Graviacypris elongata*, *Leptocythere delicata*, *L. lunata*, *L. propinqua*, *Loxococoncha endocarpa*, *Pseudostenocypris asiatica* and *Xestoleberis ementis*. *Bakunella*, *Graviacypris*, *Leptocythere*, *Loxococoncha* and *Xestoleberis* are quasi-marine forms, while *Caspiocypris* and *Pseudostenocypris* are fresh-water (see, for instance, Yassini, 1986). Four assemblage zones can be recognised on the basis of ostracods (Mamedova, in press).

The Baku Formation of Northern Iran can be correlated with the stratotypical Bakunian on ostracod evidence (Faridi, 1964; Stöcklin & Setudehnia, 1971, 1972).

Khazarian, Girkian and Khvalynian

The Khazarian takes its name from an ancient tribe who lived in the area between the Don and Volga Rivers, and the Girkian (ian) and Khvalynian take theirs from ancient names for the Caspian

Sea (Likharev, 1958). The Khazarian, Girkan and Khvalynian appear equivalent to the Uzunlarian, Karangatian and Neoeuxinian respectively of the Black Sea (Pleistocene-Holocene). Zubakov & Borzenkova (1990) calibrate the Khazarian, Girkan and Khvalynian against magnetostratigraphic polarity epoch 1 (Brunhes).

Kuyalnikian to Neoeuxinian (Black Sea) (Fig. 12)

The Kuyalnikian to Neoeuxinian of the Black Sea appear equivalent to the Akchagylia to Khvalynian of the Caspian (Late Pliocene to Holocene). Details of Kuyalnikian to Neoeuxinian stratigraphy have been discussed by Wall & Dale (1973), Karmishina (1975), Vekua (1975), Kaplin (1977), Shikmus *et al.* (1977), Schrader (1979), Semenenko & Pevzner (1979), Shatilova (1984), Sirenko & Turlo (1986), Yanko (1990a–b, 1991), Balabanov *et al.* (1991) and Trubikhin *et al.* (1991a–b).

Kuyalnikian (Fig. 12)

The Kuyalnikian takes its name from a locality on the Danube Delta (Likharev, 1958). The Romanian appears synonymous. It appears equivalent to the Akchagylia of the Caspian. Zubakov & Borzenkova (1990) calibrate it against magnetostratigraphic polarity epochs 3 (Gauss) to 2 (Matuyama).

The Kuyalnikian/Akchagylia represents a major transgressive episode when Eastern Paratethys was reconnected with the world oceans (Fig. 12). During this time, marine plankton were introduced into Eastern Paratethys enabling calibration to global datums.

Details of Kuyalnikian stratigraphy have been discussed by Karmishina (1975), Vekua (1975), Semenenko & Pevzner (1979), Shatilova (1984) and Sirenko & Turlo (1986).

Micropalaeontology. Only the non-age-diagnostic, quasi-marine, smaller benthonic foraminifera *Ammonia beccarii* and *Elphidium incertum* were recorded by Karmishina (1975) from the Kuyalnikian of Prechernomore (Northern Black Sea).

Stratigraphically and/or palaeoenvironmentally significant ostracods recorded by Karmishina (1975) and Vekua (1975) from the Kuyalnikian of Prechernomore (Northern Black Sea) and Abkhazia (North-eastern Black Sea) respectively include *Bakunella dorsoacuata*, *Caspiocypris labiata*, *Caspiolla acronasuta*, *Cryptocyprideis bogatschovi*, *Cypria arma*, *Leptocythere andrusovi*, *L. circumsulcata*, *Loxoconcha eichwaldi*, *L. petasa*, *Mediocythereis apatoica* and *Pontoniella acuminata*. *Bakunella dorsoacuata*, *Caspiolla acronasuta* and *Loxoconcha petasa* are quasi-marine species (Yassini, 1986). *Caspiocypris acronasuta* and *Cypria arma* are fresh-water.

Nannopalaeontology. Stratigraphically significant calcareous nannofossils recorded by Semenenko & Pevzner (1979) from the Kuyalnikian include *Discoaster pentaradiatus* (NN9?–NN17) and *Reticulofenestra pseudumbilica* (NN7?–NN15).

Palynology. Only non-age-diagnostic palynomorphs have been recorded from the Kuyalnikian. Pollen spectra from 'Thermo'-SCT15 are characterised by polydominance (Shatilova, 1984; Sirenko & Turlo, 1986).

Gurian

The Gurian takes its name from an ancient province of what is now Western Georgia (Likharev, 1958). It appears equivalent to

the Apscheronian of the Caspian (Pleistocene). Zubakov & Borzenkova (1990) calibrate it against magnetostratigraphic polarity epoch 2 (Matuyama).

Chaudian

The Chaudian takes its name from a promontory on the Kerch Peninsula (Crimea) (Likharev, 1958). It appears equivalent to the Bakunian of the Caspian (Pleistocene). Zubakov & Borzenkova (1990) calibrate it against magnetostratigraphic polarity epochs 2 (Matuyama) to 1 (Brunhes).

Uzunlarian, Karangatian and Neoeuxinian

The Uzunlarian takes its name from a lake, the Karangatian takes its name from a promontory on the Kerch Peninsula (Crimea), and the Neoeuxinian its name from an ancient name for the Black Sea (Likharev, 1958). The Uzunlarian, Karangatian and Neoeuxinian appear equivalent to the Khazarian, Girkan and Khvalynian respectively of the Caspian (Pleistocene-Holocene). Zubakov & Borzenkova (1990) calibrate the Uzunlarian, Karangatian and Neoeuxinian against magnetostratigraphic polarity epoch 1 (Brunhes).

Details of Uzunlarian, Karangatian and Neoeuxinian stratigraphy have been discussed by Schrader (1979) and Markova & Mikhaleska (1994). Schrader (1979) recorded the palaeoenvironmentally significant diatoms *Actinocyclus divisus*, *A. ochotensis*, *Cymatopleura solea* and *Stephanodiscus astraea* (which indicate a palaeosalinity range of 0.5–3.0ppt). Markova & Mikhaleska (1994) recorded the ostracods *Amniccythere cymbula*, *Callistocythere alfera*, *C. lopalica*, *C. mediterranea*, *C. quinquetuberculata* and *Loxoconcha immodulata* in the stratotypical Uzunlarian.

ACKNOWLEDGEMENTS. We wish to acknowledge our colleagues at the Azerbaijan Academy of Sciences, Baku, in particular Drs. Elmira Ateava, Dilara Mamedova and Reyhan Koshkarly, for their assistance in procuring publications, for facilitating fieldwork, and for sharing their extensive knowledge with us. We also wish to acknowledge Dr. Fred Rögl of the Naturhistorisches Museum, Vienna, for his constructive critical review of the manuscript. BP Exploration provided drafting facilities and is thanked for permission to publish.

REFERENCES

- Abramova, T.O. 1982. Cyclicality of spore-pollen spectra in Quaternary deposits of the west coast of the Caspian Sea. In: Kovalev, S.A. *et al.* (editors), *Sea-coasts: 32–39*. Moscow (in Russian).
- 1985. Palynological study of bottom sediments from the Caspian Sea. *Vestnik Moskovskogo Universiteta, Seriya 5 (Geologiya)*, Moscow, 1985 (5): 77–83 (in Russian).
- Adamia, Sh., Lordkipanidze, M., Beridze, M., Kotetishvili, E. & Zutelia, Z. 1990. Paleogeography of the Ukrainian Carpathians, the Crimea and the Caucasus. In: Rakus, M., Dercourt, J. & Nairn, A.E.M. (editors), *Evolution of the Northern Margin of Tethys, Vol. III: 123–146*. Paris.
- Agalarova, D.A. 1956. *Microfauna of the Productive Series and the Red-Coloured Series in Turkmenia*. Ashkhabad: Akademia Nauk Turkmen SSR (in Russian).
- 1967. *Microfauna from Pontian Sediments of Azerbaijan and adjacent regions*. Leningrad ('Nedra') (in Russian).
- Kadirova, Z.K. & Kuliyaeva, S.A. 1961. *Ostracoda in the Pliocene and Post-Pliocene deposits of Azerbaijan*. Azerbaijan Scientific Press (in Russian).
- Akrakhodzhayev, A.M., Babadagly, V.A. & Dzhumagulov, A.D. 1989. *Geology and exploration of oil- and gas-bearing ancient deltas*. Rotterdam (A.A. Balkema). English translation.
- Alekseyev, M.N. & Nikiforova, K.V. (editors). 1984. *The boundary between the*

- Neogene and Quaternary Systems in the USSR*. Moscow ('Nauka') (in Russian).
- Aleskerov, B.Dj., Mamedov, A.V., Svitch, A.A. & Yanina, T.A. 1987. New data on the stratigraphy of the Mishovdag key section in the Pleistocene of Azerbaijan. *Izvestiya Akademii Nauk Azerbajanskoi SSR, Seriya Nauk o Zemle, Baku*, 1987 (4): 50–57 (in Russian with English abstract).
- Alkhhanov, E.N. 1977. *Oil and gas in the Caspian Sea*. Moscow ('Nedra') (in Russian).
- Aliyev, Ad. A. & Aliyeva, L.I. 1970. Subdivision and correlation of Akchagylid deposits in western Azerbaijan. *Izvestiya Akademii Nauk Azerbajanskoi SSR, Baku*, 26 (2): 56–61 (in Russian).
- Aliyulla, Kh., Vekilov, B.G. & Mamedova, D.N. 1985. On the division of the stratotype section of the Baku Horizon. *Izvestiya Akademii Nauk Azerbajanskoi SSR, Seriya Nauk o Zemle, Baku*, 1985 (4): 89–93 (in Russian with English abstract).
- Ali-Zade, A.A. 1969. *Akchagylid of Azerbaijan*. Leningrad ('Nedra') (in Russian).
- 1973. *Apsheonian of Azerbaijan*. Moscow ('Nedra') (in Russian).
- 1974. The Upper Pliocene in the Caspian Sea region: ideas resulting from recent studies. *Mémoires du Bureau de Recherches Géologiques et Minières*, 2 (78): 729–731 (in French with English abstract).
- , Akhmedov, G.A., Akhmedov, A.M., Aliev, A.K. & Zeinalov, M.M. 1966. *Geology of the oil and gas fields of Azerbaijan*. Moscow (in Russian).
- & Aleskerov, J.A. 1974. Stages of Sarmatian fauna development and their role in the establishment of biostratigraphic boundaries in Azerbaijan. *Izvestiya Akademii Nauk Azerbajanskoi SSR, Seriya Nauk o Zemle, Baku*, 1974 (4): 3–8 (in Russian with English abstract).
- Ali-Zade, Ak.A., Ateava, E.Z., Koshkarly, R.O., Mamedova, D.N., Simmons, M.D. & Jones, R.W. 1994a. The biostratigraphy of the Neogene sediments of Eastern Azerbaijan and the South Caspian Basin. *Biostratigraphy of Oil and Gas Basins, St. Petersburg, Abstracts*: 7–8. VNIGNI Publication.
- , —, —, —, — & —. In press. The biostratigraphy of the Neogene sediments of Eastern Azerbaijan and the South Caspian Basin. In: Jakovleva, S. (ed.) *Biostratigraphy of Oil and Gas Basins, St. Petersburg, Proceedings VNIGNI* (in Russian).
- , Guliyev, I.S., Ateava, E.Z., Koshkarly, R.O., Mamedova, D.N., Suleymanova, S.F., Bowman, M.B.J., Brayshaw, A.C., Geehan, G., Henton, J., Jones, R.W., Reynolds, A.D., Simmons, M.D., Varney, T., Samways, G.M., Prenac, P. & Allen, L.O. 1995. Neogene stratigraphy and sedimentology in eastern Azerbaijan: outcrop observations and subsurface implications. *American Association of Petroleum Geologists, Bulletin*, 79: 1193.
- , Suleymanova, S.F., Bowman, M.B.J., Simmons, M.D., Brayshaw, A. & Reynolds, A.D. 1994b. Outcrop analogue studies and reservoir description, the Neogene Productive Series, Apsheon Peninsula, Eastern Azerbaijan. *Proceedings of the EAPG Meeting, St. Petersburg, 1994*.
- Ali-Zade, K.A. 1936. Fauna of the Akchagyl Beds of Naftalan. *Trudy Azerbajinskogo Nauchno-Issledovatel'skogo Neftyanogo Instituta*, 32 (in Russian).
- 1946. Analysis of fauna of the Productive Series. *Doklady Akademii Nauk Azerbajanskoi SSR*, 6: 3–10 (in Russian).
- 1954. *The Akchagylid Series of Azerbaijan*. Baku (in Russian).
- 1966. The stratigraphy of the Oligocene deposits of Azerbaijan. *Izvestiya Akademii Nauk Azerbajanskoi SSR, Seriya Nauk o Zemle, Baku*, 1966 (1) (in Russian).
- & Aliyeva, L.I. 1989. New data on the age of the Super-Akchagylid Continental Series of the Naftalan section, Azerbaijan. *Izvestiya Akademii Nauk Azerbajanskoi SSR, Seriya Nauk o Zemle, Baku*, 1989 (4): 121–122. (in Russian), with English abstract.
- & Atayeva, E.Z. 1982. Composition and stratigraphy of Paleogene-Miocene rocks in western Azerbaijan. *Trudy Instituta Geologii Akademii Nauk Azerbajanskoi SSR, Baku*, 1982: 60–69 (in Russian).
- , — et al. 1986. The Miocene of the south-east Caucasus and Lesser Caucasus. Pp. 222–239, in: Muratov, M.V. & Nevekskaya, L.A., *The Neogene System*. 419pp. Moscow (in Russian).
- & Mamedov, T.A. 1970. Stratigraphy of the Paleogene-Neogene boundary in Azerbaijan. *Izvestiya Akademii Nauk Azerbajanskoi SSR, Seriya Nauk o Zemle, Baku*, 1970 (3–4): 72–80 (in Russian with English abstract).
- Ananova, Ye. N. 1974. *The Neogene pollen deposits of the south of the Russian Plain*. Leningrad (Publishing House of Leningrad University) (in Russian).
- , Volkova, N.S., Zubakov, V.A., Pavlovskaya, V.I. & Remizovsky, V.N. 1985. New data on Taman key section of the Mio-Pliocene Black Sea region. *Doklady Akademii Nauk Soyuza Sotsialisticheskikh Respublik, Leningrad*, 284 (4): 925–928 (in Russian).
- Andalibi, M.J. 1991. Recent changes in Caspian Sea level. *Abstracts, XIII INQUA International Congress, Beijing, China, 1991*: 8.
- Andreescu, I. 1981. Middle–Upper Neogene and Early Quaternary chronostratigraphy from the Dacic Basin and correlations with neighbouring areas. *Annales Géologiques des Pays Helleniques hors sers.*, 4, 129–138.
- Andrusov, N.I. 1884. Geological studies in the Peninsula of Kerch in 1882 and 1883. *Notes Soc. Natur. Novoruss.*, 9 (2) (in Russian).
- Azizbekov, Sh. A. (editor). 1972. *Geology of the USSR*, vol XLVII: Azerbaijan SSR. Moscow ('Nedra') (in Russian).
- Azizbekova, I.A. 1974. Facies of Sarmatian deposits of the Nakchichevan Depression. *Izvestiya Akademii Nauk Azerbajanskoi SSR, Seriya Nauk o Zemle, Baku*, 1974 (2): 10–18 (in Russian with English abstract).
- Babayev, D.A. 1984. Features of the correlation of Pliocene deposits of uplifts of the Livanov Group and the Apsheon Archipelago. *Izvestiya Akademii Nauk Turkmenskoi SSR, Otdeleniye Fiziko-Tekhnicheskikh. Khimicheskikh i Geologicheskikh Nauk, Ashkhabad*, 1984 (5) (in Russian).
- Bagir-Zade, F.M., Narimanov, A.A. & Babayev, F.R. 1988. *Geological and geochemical features of the Caspian Sea fields*. Moscow ('Nedra').
- Balabanov, I., Yanko, V.V., Ismaylov, J. & Gey, N. 1991. The Holocene history of the Black Sea Basin. *Abstracts, XIII INQUA International Congress, Beijing, China, 1991*: 15.
- Baldi, T. 1980. The early history of Paratethys. *Bulletin of the Hungarian Geological Society*, 110, 456–472 (in Hungarian with English summary).
- Baldi, T. 1984. The terminal Eocene and Early Oligocene Events in Hungary and the separation of an anoxic, cold, Paratethys. *Eclogae Geologicae Helvetica*, 77: 1–28.
- Beard, J.H., Sangree, J.B., & Smith, L.A. 1982. Quaternary chronology, paleoclimate, depositional sequences and eustatic cycles. *American Association of Petroleum Geologists Bulletin*, 66 (2): 158–169.
- Benyamovsky, V.N., Cheltsov, Yu. G., Musikhin, V.P., Romanyuk, B.N. & Koblov, F.P. 1988. New data on the stratigraphy of the Neogene deposits of the eastern Caspian Depression. *Paleontologicheskii Sbornik, Lvov*, 25: 91–97 (in Russian).
- & Naidina, O.D. 1990. Faunal and palynological complexes of the Akchagylid of the Ural-Uil interfluvium, Eastern Precaspian. *Izvestiya Vuzov Geologii i Razvedka*, 9: 20–27 (in Russian).
- Bernor, R.L., Brunet, M., Ginsburg, L., Mein, P., Pickford, M., Rögl, F., Sen, S., Steininger, F. & Thomas, H. 1987. A consideration of some major topics concerning Old World Miocene mammalian chronology, migrations and paleogeography. *Geobios*, 20 (4): 431–439.
- , Mittmann, H.-W. & Rögl, F. 1993. Systematics and chronology of the Gotzendorf 'Hipparion' (Late Miocene, Pannonian F, Vienna Basin). *Annalen des Naturhistorischen Museums Wien*, 95 (A): 101–120.
- Blow, W.H. 1969. Late Middle Eocene to Recent planktonic foraminiferal biostratigraphy. In: Brönnimann, P. & Renz, H.H. (editors), *Proceedings of the First International Conference on Planktonic Microfossils, Geneva, 1967*: 199–412. Leiden.
- Bludorova, Ye. A., Dorofeev, L.I., Nikolaeva, K.A., Sidnev, A.V., Stepanov, L.A., Yasonov, P.G. & Yakhimovich, V.L. 1987. Lower Kama Basin. In: Alekseyev, M.N. & Nikiforova, K.V. (editors), *The boundary between the Neogene and Quaternary Systems in the USSR*: 38–44. Moscow ('Nauka') (in Russian).
- Bogdanowicz, A.K. 1947. Results of the investigations of Miocene foraminifera from the Crimea and Caucasus. *Trudy Vsesoyuznogo Neftyanogo Nauchno-Issledovatel'skogo Geologo-Razvedochnogo Instituta (VNIGRI), Leningrad & Moscow*, 1947: 5–33 (in Russian).
- 1950a. Stratigraphical subdivision of the Tarkhanian of Kuban on the basis of foraminiferal investigations. *Trudy Vsesoyuznogo Neftyanogo Nauchno-Issledovatel'skogo Geologo-Razvedochnogo Instituta (VNIGRI), Novaya Seriya, Leningrad & Moscow*, 51: 113–128 (in Russian).
- 1950b. Tschokrakian Foraminifera of the western Precaucasus. *Trudy Vsesoyuznogo Neftyanogo Nauchno-Issledovatel'skogo Geologo-Razvedochnogo Instituta (VNIGRI), Novaya Seriya, Leningrad & Moscow*, 51: 129–176 (in Russian).
- 1965. New data on Konkian Miliolidae from the western Precaucasus. *Trudy Vsesoyuznogo Neftyanogo Nauchno-Issledovatel'skogo Geologo-Razvedochnogo Instituta (VNIGRI), Paleontologicheskii Sbornik 2, Leningrad & Moscow*, 16: 34–49 (in Russian).
- 1967. New types of Articuline from the Maeotian of Kuban. *Paleontologicheskii Zhurnal, Moscow*, 1: 131–132 (in Russian).
- 1969. Maeotian Miliolidae from the western Precaucasus. *Trudy Vsesoyuznogo Neftyanogo Nauchno-Issledovatel'skogo Geologo-Razvedochnogo Instituta (VNIGRI), Paleontologicheskii Sbornik 2, Leningrad & Moscow*, 19: 64–133 (in Russian).
- 1974. New Miliolids from the Meotian in Western Georgia. *Trudy Vsesoyuznyi Nauchno-Issledovatel'skii Geologorazvedochnyi Neftyanoi Institut (VNIGNI), Leningrad*, 152: 115–125.
- Bolli, H.M. & Krascheninnikov, V.A. 1977. Problems in Paleogene and Neogene correlations based on planktonic foraminifera. *Micropaleontology*, 23 (4): 436–452.
- Boomer, I. 1993a. Palaeoenvironmental indicators from Late Holocene and contemporary Ostracoda of the Aral Sea. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 103: 141–153.
- 1993b. Sub-fossil Ostracoda and the death of the Aral Sea. *Geology Today (Jan.–Feb. 1993)*: 18–22.
- Brzobohaty, R. 1983. Fish otoliths from the West Carpathian Tertiary and their biostratigraphic significance. In: Thon, A. (editor), *Miscellanea*

- Micropalaeontology*: 247–266. Hodonin.
- Bugrova, E.M. 1986. Detail of biostratigraphic division of the Eocene of the Krasnovodsk Peninsula and the Karabogaz'ya region according to foraminifera. In: Krymgo'ts, G. Ya. & Belenkova, V.S. (editors), *Paleontology and Detailed Stratigraphic Correlation*: 41–49. Leningrad ('Nauka') (in Russian).
- Bukry, D. 1974. Coccoliths as paleosalinity indicators – evidence from the Black Sea. In: Degens, E.T. & Ross, D.A. (editors), *The Black Sea – Geology, Geochemistry and Biology*: 353–363. Tulsa.
- Buryakovskiy, L.A. 1974. Distribution patterns of oil and gas deposits within the Apsheron Archipelago. *International Geological Review*, 16 (7): 749–758.
- 1993. Outline of general and petroleum geology in Azerbaijan and the South Caspian Basin. *Bulletin of the Houston Geological Society*, February 1993: 16–47.
- Chelidze, A.L. 1973. Correlation of the Pontian from the Euxinian and Caspian Basins. *Soobshcheniya Akademii Nauk Gruzinskoi SSR, Tbilisi*, 71 (3): 645–648 (in Russian with English abstract).
- Chepal'ga, A.L. 1985. Climatic and eustatic fluctuations in the Paratethys Basin's history. *Abstracts, VII Congress, Regional Commission on Mediterranean Neogene Stratigraphy (RCMNS)*, Budapest, 1985: 134–136.
- 1991. Pleistocene Black Sea level changes and ecological consequences. *Abstracts, XIII INQUA International Congress, Beijing, China, 1991*: 56.
- , Korotkevich, E.L., Trubikhin, V.M. & Svellitskaya, T.V. 1985. Chronology of the Eastern Paratethys regional stages and Hipparion events according to palaeomagnetic data. *Abstracts, VII Congress, Regional Commission on Mediterranean Neogene Stratigraphy (RCMNS)*, Budapest, 1985: 137–139.
- Chumakov, I.S., Byzova, S.L. & Ganzey, S.S. 1988. Maotian to Pontian geochronology of Eastern Paratethys. *Doklady Akademii Nauk SSSR*, 303 (1): 178–181 (in Russian).
- , — & — 1992a. *Cenozoic geochronology and correlation of Eastern Paratethys*. Moscow ('Nauka') (in Russian).
- , —, Arias, C., Bigazzi, G., Bonadonna, F.P., Hadler Neto, J.C. & Norelli, P. 1992b. Interlaboratory fission track dating of volcanic ash levels from Eastern Paratethys: a Mediterranean-Paratethys correlation. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 95: 287–295.
- , Ganzey, S.S., Byzova, S.L., Dobrynya, V.Ya. & Paramonova, N.P. 1984. Sarmatian geochronology of Eastern Paratethys. *Doklady Akademii Nauk SSSR*, 276 (5): 1189–1193 (in Russian).
- Cicha, I., Marinescu, F. & Senes, J. 1975. *Correlation du Neogene de la Paratethys Centrale*. Prague.
- , Senes, J. & Tejkal, J. 1967. *Chronostratigraphie und Neostratotypen Miozen der Zentralen Paratethys*. I. Karpatien. Bratislava.
- , Zapletalova, I., Molchikova, V. & Brzobohaty, R. 1983. Stratigraphical range of Eggenburgian-Badenian Foraminifera in West Carpathian Basins. In: Thon, A. (editor), *Miscellanea Micropalaeontologia*: 99–144. Hodonin.
- de Deckker, P. 1981. Ostracods of athalassic saline lakes. *Hydrobiologia*, 81: 131–144.
- Demarcq, J. 1985. Paleothermic evolution during the Neogene in Mediterranean through the marine megafauna. *Abstracts, VII Congress, Regional Commission on Mediterranean Neogene Stratigraphy (RCMNS)*, Budapest, 1985: 176–178.
- Dercourt, J., Ricou, L.E., Adamia, Sh., Csaszar, G., Funk, H., Lefelf, J., Rakus, M., Sandulescu, M., Tollmann-A. & Tchoumachenko, P. 1990. Anisian to Oligocene paleogeography of the European margin of Tethys (Geneva to Baku). In: Rakus, M., Dercourt, J. & Nairn, A.E.M. (editors), *Evolution of the Northern Margin of Tethys*, Vol. III: 159–190. Paris.
- Dmitriyev, A.V., Nazarov, N.O., Nigarov, A. & Uzakov, O. 1969. Studies on the Akchaglyan of Turkmenia. *Izvestiya Akademii Nauk Turkmenkoi SSR, Seriya Fiziko-Tekhnicheskikh, Khimicheskikh i Geologicheskikh Nauk, Ashkhabad*, 1969 (3): 125–127 (in Russian).
- Dumitrica, P. 1985. Silicoflagellates – biostratigraphic and paleogeographic markers for Middle Miocene of Central Paratethys. *Abstracts, VII Congress Regional Commission on Mediterranean Neogene Stratigraphy (RCMNS)*, Budapest, 1985: 188–190.
- Dzhabarova, Kh.S. 1966. The lithostratigraphic and geochemical characteristics of the Akchagly deposits in the eastern part of the Kura Basin. *Izvestiya Akademii Nauk Azerbajdzhanskoi SSR, Seriya Nauk o Zemle*, 1966 (5): 3–7 (in Russian).
- 1973. Stratigraphic subdivision of Middle Miocene deposits of Middle Kura Depression on palynological data. *Palynologiya Kaynofta, Moscow*, 1973: 173–177 (in Russian with English abstract).
- 1978. Development of Neogene flora of Azerbaijan on the basis of palynological data. *Izvestiya Akademii Nauk Azerbajdzhanskoi SSR, Seriya Nauk o Zemle, Baku*, 1978 (5): 60–65 (in Russian with English abstract).
- 1980. Developmental stages of the Upper Palaeogene and Neogene floras of Azerbaijan according to palynological data. *Proceedings of the Fourth International Palynological Conference, Lucknow, 1975–1976*, 2: 747–749.
- Dzhanelidze, O. I. 1970. *Foraminifera of the Lower and Middle Miocene of Georgia*. Tbilisi: 'Metsnerieva' (in Russian).
- 1977. *Polymorphinids of the Miocene of Georgia*. Tbilisi: 'Metsnerieva' (in Russian).
- , Vekua, M.L. & Maizuradze, L.S. 1985. *Development of Ostracod and Foraminifera Fauna of the Neogene Black Sea/Caspian Sea Zone*. Tbilisi: 'Metsnerieva' (in Russian).
- Faridi, Z. 1964. Ostracods in the Plio-Pleistocene sediments of Gorgan and Mazanderan (Northern Iran). *Iran Petroleum Institute Bulletin*, 14: 534–535.
- Fedkovich, Z.N. 1978. Comparative analysis of complexes of Akchaglyan mollusks in the Northern Caspian Depression, the Southern Volga region near Kuibyshev and the Uralian Foredeep near Drenburg, and at the same time in Turkmenia and Azerbaijan. *Voprosy Paleontologii i Stratigrafii Azerbajdjanu, Baku*, 1978: 51–61 (in Russian).
- Fedorov, P.V. 1994. Some problems concerning the geological history of the Caspian Sea. *Stratigraphy and Geological Correlation*, 2 (2): 71–79.
- Gasanol, T.A. & Kyazymov, T.M. 1988. Paleogene and Neogene sediment accumulation in eastern Lesser Caucasus. *Soviet Geology, Moscow*, 1988 (3): 62–70.
- Gasanova, M.S. 1965. *Diatomaceous flora of the Middle and Upper Miocene of Kobustan (eastern Azerbaijan)*. First All-Union Palaeontological Council, Novosibirsk (in Russian).
- Ghanbari, E. 1991. Neogene-Quaternary tectonics in Azerbaijan area (Iran). *Abstracts, XIII INQUA International Congress, Beijing, China, 1991*: 47.
- Gheorghian, M. 1974. Distribution pattern of benthonic foraminifera on continental shelf of Black Sea off Romanian shore. In: Degens, E.T. & Ross, D.A. (editors), *The Black Sea – geology, geochemistry and biology*: 411–419. Tulsa.
- Gofman, A.A. 1966. *Ecology of the contemporary and Novocaspian ostracods of the Caspian Sea*. Moscow ('Nauka') (in Russian).
- Gramann, F. 1971. Brackish or hyperhaline? Notes on paleoecology based on Ostracoda. *Bulletin du Centre de Recherche de Pau*, 5: 93–99.
- Grichuk, V.P. (editor). 1973. *Pliocene and Pleistocene palynology*. Moscow ('Nauka') (in Russian).
- 1984. Late Pleistocene vegetation history. In: Velichko, A.A. (editor), *Late Quaternary environments of the Soviet Union*: 155–178. London (Longman) (English translation).
- , Gurtovaya, Ye. Ye., Zelikson, E.M. & Borisova, O.K. 1984. Methods and results of Late Pleistocene paleoclimatic reconstruction. In: Velichko, A.A. (editor), *Late Quaternary Environments of the Soviet Union*: 251–260. London (Longman) (English translation).
- Grishanov, A.N., Yereimin, V.N., Imnadze, Z.A., Kitovani, T.G., Molotovskiy, E.A. & Torozov, R.I. 1983. Stratigraphy of Upper Pliocene and Lower Pleistocene of Fourii (western Georgia) from paleontology and paleomagnetism. *Byulleten' Komissii po Izucheniyu Chetvertichnogo Perioda*, 52: 18–28 (in Russian).
- Haq, B.U., Hardenbol, J. & Vail, P.R. 1988. Mesozoic and Cenozoic chronostratigraphy and eustatic cycles. In: Wilgus, C.K. et al. (editors), *Sea-level changes: an integrated approach*: 71–108. Society of Economic Paleontologists and Mineralogists (Special Publication, No. 42).
- Hsu, K.J. 1983. *The Mediterranean was a desert: a voyage of the Glomar Challenger*. Princeton (University Press).
- Imnadze, Z.A. 1964. Some data on the ostracods of the Pliocene deposits of western Georgia. *Voprosy Geologii Gruzii, Acad. Nauk. Gruzinski SSR, Geol. Inst., XXII Sess.*, 365–371 (in Russian).
- 1974. Ostracods of the Pliocene of Western Georgia. *Trudy Vsesoyuznyi Nauchno-Issledovatel'skii Geologorazvedochnyi Neftyanoi Institut (VNIGNI), Leningrad*, 152: 126–133.
- & Karmishina, G.I. 1980. Comparison of Pliocene Ostracoda complexes in the northern and eastern Black Sea Region. *Voprosy Stratigrafii, Leningrad*, 5: 131–148 (in Russian).
- Iossofova, Yu. I. 1985. The Neogene history of the Palaeo-Don Basin. *Abstracts, VII Congress Regional Commission on Mediterranean Neogene Stratigraphy (RCMNS)*, Budapest, 1985: 271–273.
- Ismailov, K.A. & Idrisov, V.G. 1963. *Oil and gas deposits of the eastern part of the Apsheron Peninsula*. Baku (Inst. Akad. Nauk Azerbaijan SSR).
- , — & Tagiev, E.A. 1972. *The Productive Series of the Baku Archipelago*. Baku ('Elm').
- Ivanova, N.G. 1985. Palynological evaluation of Apsheron deposits of western Turkmenia. In: Grichuk, V.P. & Zaklinskaya, E.D. (editors), *Palynology of the Quaternary Period*: 158–167. Moscow ('Nauka') (in Russian).
- Jiricek, R. 1975. *Biozonen der Zentralen Paratethys*. Gbely, CSSR (NAFTA).
- 1983. Redefinition of the Oligocene and Neogene Ostracod zonation of the Paratethys. In: Thon, A. (editor) *Miscellanea Micropalaeontologia*: 195–236. Hodonin.
- Jones, R.W. 1996. *Micropalaeontology in Petroleum Exploration*. Oxford University Press.
- Kaplin, P.A. (editor). 1977. *Paleogeography and deposits of the Pleistocene of the southern seas of the USSR*. Moscow ('Nauka') (in Russian).
- Karmishina, G.I. 1975. *Pliocene Ostracoda from the southern (European) USSR*. Saratov (Izd. Saratovskogo Univ.) (in Russian).
- Keraudren, B. & Thibault, C. 1987. On the Plio-Pleistocene formations of the Iranian sector of the Caspian Sea. *Proceedings, International Symposium on*

- Shallow Tethys 2, Wagga Wagga*: 141–149 (in French).
- Kerimov, V.U., Khalilov, E.A. & Mekhtiev, N.U. 1991. The paleogeographic conditions in the formation of the South Caspian Depression in the Pliocene in connection with its gas- and oil-bearing capacity. *Geologia Nefti y Gaza*, **10**: 5–8 (in Russian with English abstract).
- Khain, V., Apressov, S. & Nirtchink, M. 1937. *Petroleum excursion notes (Azerbaijani Soviet Socialist Republic)*. Moscow, Leningrad.
- Khalilov, D.M. 1946. Microfauna of the Productive Series of Azerbaijan. *Doklady Akademii Nauk Azerbajanskoi SSR*, **6**: 11–21 (in Russian).
- & Mamedova, L.D. 1973. Zonal subdivision of Paleogene sediments in Azerbaijan based on foraminifera. *Ocherki po Geologii Azerbajanskoi SSR, Baku*, 1973: 218–226 (in Russian).
- Khotinskiy, N.A. 1984. Holocene climatic change. In: Velichko, A.A. (editor), *Late Quaternary Environments of the Soviet Union*: 179–200. London (Longman) (English translation).
- Kleschev, K.A., Shein, V.S. & Clavkin, V.S. 1992. New concepts of geological structure and oil and gas presence in western Turkmenia. *Geologiya Nefti i Gaza*, **5**: 2–6 (in Russian).
- Klopotskaya, N.B. 1991. The Caucasus at the coldest time of the last glaciation. *Abstracts, XIII INQUA International Congress, Beijing, China, 1991*: 164–165.
- Knystautas, A. 1987. *The natural history of the Soviet Union*. London (Century).
- Kopp, L.M. 1970. Upper Pliocene stratigraphy in the western branches of the Kopet Dag. *Izvestiya Vysshikh Uchebnykh Zavedenii, Geologiya i Razvedka, Moscow*, **6**: 14–20 (in Russian).
- Korenova, E.V. & Kartashova, G.G. 1978. Palynological study of samples from Holes 379A, 380A, Leg 42B. *Initial Reports of the Deep Sea Drilling Project*, **42B**: 951–992.
- Koshkarly, R.O. 1986. Features of nannoplankton distribution in the Paleogene deposits of Azerbaijan. *Izvestiya Akademii Nauk Azerbajanskoi SSR, Seriya Nauk o Zemle, Baku*, **1986** (1): 107–110 (in Russian with English abstract).
- 1993. Calcareous nannoplankton biostratigraphy of the Palaeogene deposits of Azerbaijan. *Knikhovnicka ZPN*, **14B** (2): 63–68.
- & Alekperov, N. 1993. Biostratigraphy of the Paleogene deposits of Azerbaijan by nannoplankton. *Abstracts, Geological Congress of Turkey, 1993*: 50.
- & Baldi-Beke, M. 1987. Correlation of the Paleogene deposits of Azerbaijan and Hungary by nannoplankton. *Acta Geologica Hungarica*, **30** (3–4): 289–298.
- Krakhmalnaya, T.V., Svetlitskaya, T.V. & Chepalyga, A.L. 1993. New data on stratigraphy, magnetostratigraphy and mammal faunas of the late Miocene locality of Novaya Emetovka (Ukraine). *Newsletters on Stratigraphy*, **29**: 77–89.
- Krasheninnikov, V.A. 1986. The Kuban River sequence (USSR, North Caucasus). In: Pomerol, Ch. & Premoli Silva, I. (editors), *Terminal Eocene Events*: 137–139. Amsterdam (Elsevier).
- & Muzylev, N.G. 1975. Relationship between the zonal scales based on planktonic foraminifera and nannoplankton in Paleogene sections of the north Caucasus. *Voprosy Mikropaleontologii, Moscow*, **1975**: 212–224 (in Russian with English abstract).
- & Ptukhina, A.E. 1985. Stratigraphical subdivision of Paleogene deposits of Armenia by planktonic foraminifera, nannoplankton and nummulites (Pt. 1, reference Paleogene sections of Armenia). *Voprosy Mikropaleontologii, Moscow*, **1985**: 130–169 (in Russian with English abstract).
- & Ptukhina, A.E. 1986. Stratigraphical subdivision of Armenian Paleogene deposits by planktonic microfossils and nummulites (regional stratigraphy, zonal scales by planktonic and benthonic microfossils, their correlation). *Voprosy Mikropaleontologii, Moscow*, **1986**: 60–98 (in Russian with English abstract).
- Kretzoi, M. 1985. Sketch of the biochronology of the Late Cenozoic in Central Europe. Pps. 3–20, in Kretzoi, M. & Pecsli, M. (editors): *Problems of the Neogene and Quaternary in the Carpathian Basin*. Budapest (Akademiai Kiado).
- Krhovsky, J. 1985. Central Paratethys ecostratigraphic correlations in relation to the Oligocene sea-level changes. *Abstracts, VII Congress Regional Commission on Mediterranean Neogene Stratigraphy (RCMNS)*, Budapest, 1985: 333–335.
- Krstic, N. 1976. Pontian ostracodes in Paratethys and Tethys. *Proceedings, VI Congress Regional Commission on Mediterranean Neogene Stratigraphy (RCMNS)*, Bratislava, 1975: 325–330.
- Kuliyeva, S.A. 1968. Stratigraphy of the Apsheron Suite in the Baku Archipelago according to microfaunas. *Doklady Akademii Nauk Azerbajanskoi SSR, Baku*, **24** (8): 52–54 (in Russian).
- Lamb, J.L., Wornardt, W.W., Huang, T.-C. & Dube, T.E. 1987. Practical application of Pleistocene eustasy in offshore Gulf of Mexico. *Special Publications of the Cushman Foundation for Foraminiferal Research*, **24**: 33–40.
- Likharev, B.K. (Dir.). 1958. *U.R.S.S. Lexique stratigraphique internationale*, Paris, II (1–4).
- Lindsay, E.H., Fahlbusch, V. & Mein, P. 1989. *European Neogene mammal chronology*. Plenum Press.
- Livental, V.E. 1929. Ostracods from the Akchagyl and Apsheronian Beds of the Babazan Section. Baku (Izd. Rabot Issled. Lab. Po Geol. Neft) (in Russian).
- Lupov, N.P., Semenov, V.V. & Smirnov, L.N. (editors). 1972. *Geology of the USSR, Vol. XXII: Turkmenian SSR (Geological Description)*. Moscow ('Nedra') (in Russian).
- Luttig, G. & Steffens, P. 1976. *Paleogeographic atlas of Turkey from the Oligocene to the Pleistocene*. Hanover.
- Macarovic, N. & Cehan-Jones, B. 1962. Distribution des foraminifères sur la plate-forme continentale du nord-ouest de la Mer Noire. *Travaux du Musée Histoire Naturelle 'Gr. Antipa'*, Bucharest, **3**: 45–60.
- Maisuradze, P.C. 1971. Sarmatian Foraminifera of Western Georgia. Tbilisi, 'Metsnerieva' (in Russian).
- 1988. *Foraminifera of the Maeotian of West Georgia*. Tbilisi: 'Metsnerieva' (in Russian).
- Malaeva, E.V. & Kulikov, O.A. 1991. Late Pleistocene pollen floras from the dated loess sequence of Kopetdag (Turkmenia). *Abstracts, XIII INQUA International Congress, Beijing, China, 1991*: 229.
- Mamedov, A.V. & Aleskerov, B.D. 1985. Stratigraphy, chronology and paleogeography of the Pleistocene in Azerbaijan and the Caspian regions. *Izvestiya Akademii Nauk Azerbajanskoi SSR, Seriya Nauk o Zemle, Baku*, **1985** (3): 46–54 (in Russian with English abstract).
- & — 1986. Stratigraphy, chronology and paleogeography of the Pleistocene in Azerbaijan. *Izvestiya Akademii Nauk SSSR, Seriya Geologicheskaya, Leningrad*, **1986** (1): 93–104 (in Russian).
- & Rabotina, Y.N. 1984a. Reconstruction of some climatic elements of the Upper Pliocene in Azerbaijan. *Izvestiya Akademii Nauk Azerbajanskoi SSR, Seriya Nauk o Zemle, Baku*, **1984** (3): 15–22 (in Russian with English abstract).
- & — 1984b. The vegetation of Azerbaijan in the Early and Middle Pleistocene. *Izvestiya Akademii Nauk Azerbajanskoi SSR, Seriya Nauk o Zemle, Baku*, **1984** (6): 10–16 (in Russian with English abstract).
- Mamedov, P.Z. 1989. Paleo-deltaic complexes in the north of the South Caspian Depression. *Petroleum Geology*, **25** (9/10): 344–346.
- Mamedova, D.N. 1984. The change of ostracod assemblages at the Apsheron/Baku boundary (materials of the Baku Horizon stratotype). *Izvestiya Akademii Nauk Azerbajanskoi SSR*, **1984** (2): 127–130 (in Russian, with English abstract).
- 1985. Ostracods of the type Baku Stage. *Dep. VINITI*, **6**: 1–4 (in Russian).
- 1988. On the study of Ostracoda of the Apsheronian Stage in the Yasamal Valley. *Izvestiya Akademii Nauk Azerbajanskoi SSR*, **1988** (4): 53–57 (in Russian, with English summary).
- 1993. Ostracodes of the stratotype of Lower Pleistocene in the eastern Caucasus. *Abstracts, Geological Congress of Turkey, 1993*: 49.
- In press. Ostracod biostratigraphy of the Pliocene – Pleistocene sediments of eastern Azerbaijan. In: Simmons, M.D. (ed.) *Recent Developments in Micropaleontology in the Former Soviet Union*. Chapman & Hall.
- Mamedova, L.D. 1971. On the stratigraphy of Miocene deposits of north-eastern Azerbaijan in the light of the study of its microfauna. *Doklady Akademii Nauk Azerbajanskoi SSR, Baku*, **27** (7): 37–41 (in Russian with English abstract).
- 1987. The sequence of Miocene deposits in the Turzhanchay River valley. *Doklady Akademii Nauk Azerbajanskoi SSR, Baku*, **43** (6): 79–82 (in Russian with English abstract).
- & Mamedova, N.A. 1970. Paleogene stratigraphy near the village of Gadzhilu in the Shemakha region, Azerbaijan. Based on Foraminiferal and Radiolarian Faunas. *Doklady Akademii Nauk Azerbajanskoi SSR, Baku*, **26** (9): 53–56 (in Russian).
- Mandelstam, M.I., Markova, P., Rozyeva, T.R. & Stepanaitas, N.E. 1962. *Ostracods from Pliocene and Post-Pliocene sediments of Turkmenistan*. Ashkhabad (Izd. Akad. Nauk. Turkmen SSR) (in Russian).
- Markova, A.K. & Mikhailesku, K.D. 1994. Correlation of Pleistocene marine and continental deposits from the Northwestern Black Sea region. *Stratigraphy and Geological Correlation*, **2** (4): 388–394.
- Martini, E. 1971. Standard Tertiary and Quaternary calcareous nannoplankton zonations. In: Farinacci, A. (editor), *Proceedings of the Second International Conference on Planktonic Microfossils, Rome, 1969*: 739–785. Rome (Edizioni Tecnoscienza).
- Mekhtiev, Sh.F. & Pashaly, N.V. 1987. Lithology of Middle Pliocene Deposits of the Southern Caspian and their correlation. In: *Lithology and Mineral Resources*: 56–72. Plenum Publishing Corporation.
- Meszaros, N. 1992. Nannoplankton zones in the Miocene deposits of the Transylvanian Basin. *International Nannoplankton Association Newsletter*, **13** (2): 59–60.
- Muratov, M.V. 1960. *A brief summary of the geological structure of the Crimean Peninsula*. Moscow (in Russian).
- & Nevesskaya, P.A. 1986. *Neogene System*. Moscow; Palaeontological Institute (in Russian).
- Murray, J.W. 1973. *Distribution and ecology of living benthic foraminifera*. London (Heinemann Educational Books).
- 1991. *Ecology and palaeoecology of benthic foraminifera*. London (Longman Scientific & Technical).
- Muzylev, N.G. & Golovina, L.A. 1987. The link of the Eastern Paratethys and the world ocean in the early Middle Miocene. *Izvestiya Akademii Nauk SSSR, Seriya Geologicheskaya*, **12**: 67–73 (in Russian).

- [illegible]

- & Steininger, F.F. 1983. Vom Zerfall der Tethys zu Mediterran und Paratethys. *Annalen des Naturhistorischen Museums Wien*, **85/A**: 135–163.
- & — 1984. Neogene Paratethys, Mediterranean and Indo-Pacific seaways. In: Brencley, P. (editor), *Fossils and Climate*: 171–200. London.
- , — & Muller, C. 1978. Middle Miocene salinity crisis and paleogeography of the Paratethys (middle and eastern Europe). In: Ross, D.A., Neprochnov, Y.P. et al. (editors), *Initial Reports of the Deep Sea Drilling Project*, **42**: 985–990.
- , Zapfe, H., Bernor, R.L., Brzobohaty, R.L., Daxner-Hock, G., Draxler, I., Fejfar, O., Gaudant, J., Herrmann, P., Rabeder, G., Schultz, O. & Zetter, R. 1993. Die Primatenfundstelle Gotzendorf an der Leitha (Obermiozän des Wiener Beckens, Niederösterreich). *Jahrbuch der Geologischen Bundesanstalt*, **136** (2): 503–526.
- Ross, D.A., Neprochnov, Y.P. et al. (editors). 1978. *Initial Reports of the Deep Sea Drilling Project*, **42**.
- Royden, L.H. & Horvath, F. (editors). 1987. *The Pannonian Basin: a study in basin evolution*. Tulsa, Oklahoma (American Association of Petroleum Geologists, Memoir No. 45).
- Rozyeva, T.R. 1971. Composition and stratigraphic distribution of the Tertiary ostracods of Turkmenia. In: Vyalov, O.S. (editor), *Fossil Ostracoda*: 169–178. Jerusalem (Israel Program for Scientific Translation).
- Rusu, A. 1985. Oligocene events in Transylvania and the first separation of Paratethys. *Abstracts, VII Congress Regional Commission on Mediterranean Neogene Stratigraphy*, Budapest, 1985: 502.
- Savina, S.S. & Khotinskiy, N.A. 1984. Holocene paleoclimatic reconstruction based on the zonal method. In: Velichko, A.A. (editor), *Late Quaternary Environments of the Soviet Union*: 287–296. London (Longman) (English translation).
- Schrader, H.-J. 1979. Quaternary paleoclimatology of the Black Sea. *Sedimentary Geology*, **23**: 165–180.
- Semenenko, V.N. 1979. Correlation of Mio-Pliocene of the Eastern Paratethys and Tethys. *Annales Géologiques des Pays Helléniques*, **1979**: 1101–1111.
- 1984. *Stratigraphic correlation of the Upper Miocene and Pliocene of the Eastern Paratethys and Tethys (the Mediterranean)*. Moscow (State University Dissertation) (in Russian).
- & Lulieva, S.A. 1982. The problems of direct correlation in the Upper Miocene and Pliocene of the Eastern Paratethys and Tethys. *Izvestiya Akademii Nauk SSSR, Seriya Geologicheskaya*, **9**: 61–71 (in Russian).
- & — 1985. About the Miocene-Pliocene boundary in Paratethys. *Abstracts, VII Congress Regional Commission on Mediterranean Neogene Stratigraphy (RCMNS)*, Budapest, 1985: 507–508.
- & Pevzner, M.A. 1979. Biostratigraphic and paleomagnetic correlation of the Miocene and Pliocene in the Pontic-Caspian region. *Izvestiya Akademii Nauk SSSR, Seriya Geologicheskaya*, **1979** (1): 5–15 (in Russian).
- Senes, J. 1973. Einleitung in Chronostratigraphie und Neostatotypen Miozän der Zentralen Paratethys. Verlag Slowakischen Akademie Wissenschaftlichen, Bratislava, **3**: 21–25.
- 1985a. The stratigraphic correlation of the Tethys-Paratethys Neogene. *Abstracts, VII Congress Regional Commission on Mediterranean Neogene Stratigraphy (RCMNS)*, Budapest, 1985: 33–42.
- 1985b. Stratigraphic correlation Tethys-Paratethys Neogene: IGCP Project No. 25 (history and aims, organization, results). *Geologicky Zbornik, Bratislava*, **36** (6): 725–746.
- & Marinescu, F. 1974. Cartes paléogéographiques du Neogene de la Paratethys. *Mémoires du Bureau de Recherches Géologiques et Minières*, **88** (2): 767–774.
- Shatlova, I.I. 1984. Late Miocene vegetation of western Georgia. Tbilisi ('Metsniereva') (in Russian).
- Shchekina, N.A. 1979. The history of flora and vegetation in the south of the European part of the USSR during the Late Miocene-Early Pliocene. Kiev ('Naukova Dumka') (in Russian).
- Sheydayeva-Kuliyeva, Kh. M. 1966. *Ostracoda in the Pontian layer of Azerbaijan*. Ashkabad (Izd. Akad. Nauk Turkmen SSR) (in Russian).
- Shihliniski, A.S. 1967. *Geology and prospectivity of hydrocarbon deposits of the Pliocene of the Lower Kura Basin*. Baku.
- Shikmuss, K.M., Komarov, A.V. & Grakova, I.V. 1977. Stratigraphy of the Upper Quaternary deep-sea sediments in the Black Sea. *Oceanology*, **17** (4): 443–446.
- Shishova, Z.A. 1955. New data from the study of diatomaceous algae in the Miocene of the Apsheron Peninsula. *Doklady Akademii Nauk Azerbajjanskoi SSR, Baku*, **11** (6) (in Russian).
- Simmons, M.D., Ali-Zade, Ak.A., Allen, L.O., Ateava, E.Z., Baba-Zade, A.A., Brenac, P., Jones, R.W., Koshkarily, R.O., Mamedova, D.N. & Suleymanova, S.F. In Press. Biostratigraphy of the Neogene Succession in Eastern Azerbaijan. In: Simmons, M.D. (ed.) *Recent Developments in Micropalaeontology in the Former Soviet Union*. Chapman & Hall.
- Sirenko, N.A. & Turlo, S.I. 1986. *Development of the Ukrainian soil and vegetation through the Pliocene and Pleistocene*. Kiev ('Naukova Dumka') (in Russian).
- Skalbdyna, L.N. (editor). 1985. *Geological events in the history of the Pliocene and Pleistocene of southern and northern seas*. Moscow (Akad. Nauk SSSR) (in Russian).
- Slama, P. 1983. Lower Badenian Radiolaria in the Moravian part of the Carpathian Foredeep. In: Thon, A. (editor), *Miscellanea Micropalaeontologia*: 145–170. Hodonin.
- Spiegler, D. von & Rögl, F. 1992. *Bolboforma* (Protohyta, Incertae Sedis) im Oligozän und Miozän des Mediterrän und der Zentralen Paratethys. *Annalen des Naturhistorischen Museums Wien*, **94** (A): 59–95.
- Steininger, F.F. & Papp, A. 1979. Current biostratigraphic and radiometric correlations of Late Miocene Central Paratethys stages (Sarmatian s.str., Pannonian s.str. and Pontian) and Mediterranean stages (Tortonian and Messinian) and the Messinian event in the Paratethys. *Newsletters on Stratigraphy*, **8** (2): 100–110.
- , Rabeder, G. & Rögl, F. 1985. Land mammal distribution in the Mediterranean Neogene: a consequence of geokinematic and climatic events. In: Stanley, D.J. & Wezel, F.-C. (editors), *Geological Evolution of the Mediterranean Basin*: 559–571. New York (Springer-Verlag).
- & Rögl, F. 1979. The Paratethys history – a contribution towards the Neogene dynamics of the Alpine orogen (an abstract). *Annales Géologiques des Pays Helléniques*, **1979**: 1153–1165.
- & — 1984. Paleogeography and palinspastic reconstructions of the Neogene of the Mediterranean and Paratethys. In: Dixon, J.E. & Robertson, A.H.F. (editors), *The Geological Evolution of the Eastern Mediterranean*: 659–668. London. (4–15–10–11)
- , —, Hochuli, P. & Muller, C. 1989. Lignite deposition and marine cycles: the Austrian Tertiary lignite deposits – a case history. *Sitzungsberichte der Österreichischen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse, Abt. I*, **197** (5–10): 309–332.
- , — & Martini, E. 1976. Current Oligocene/Miocene biostratigraphic concept of the Central Paratethys (middle Europe). *Newsletters on Stratigraphy*, **4** (3): 174–202.
- , Senes, J., Kleeman, K. & Rögl, F. (editors). 1985. *Neogene of the Mediterranean Tethys and Paratethys: stratigraphic correlation tables and sediment distribution Maps*. Vienna (University Press). 1071 10 1011 102027
- Stöcklin, J. & Setudehnia, A.O. 1971. *Stratigraphic Lexicon of Iran (Part I – Central, North and East Iran; Part II – Southwest Iran)*. Tehran; Geological Survey of Iran (Report, 18).
- & — 1972. *Iran. Lexique Stratigraphique International*, **III** (9b).
- Subbotina, N.N., Pishvanova, L.S. & Ivanova, L.V. 1960. Stratigraphy of the Oligocene and Miocene deposits of the Pre-Carpathians according to the foraminifera. *Trudy Vsesoyuznogo Neftyanago Nauchno-Issledovatel'skogo Geologo-Razvedochjnogo Instituta (VNIGRI), Leningrad & Moscow*, **153**: 5–146 (in Russian).
- Sultanov, A.D. 1979. *Lithology of Akchagyljan deposits of Azerbaijan*. Baku ('Elm') (in Russian).
- & Sheydayeva-Kuliyeva, Kh. M. 1969. On the lithofacial and faunistic characteristics of the Akchagyljan deposits of Azerbaijan. *Doklady Akademii Nauk Azerbajjanskoi SSR, Baku*, **25** (10): 47–49 (in Russian).
- Sultanov, E.T. 1985. Paleogeography of the Early and Middle Pleistocene in the southeastern Caucasus. *Izvestiya Akademii Nauk Azerbajjanskoi SSR, Seriya Nauk o Zemle, Baku*, **1985** (6): 96–102 (in Russian with English abstract).
- Sultanov, K.M. 1964. *The Apsheron Stage in Azerbaijan*. Baku (in Russian).
- Suzin, A.V. 1956. *Ostracods of the Tertiary of the northern Pre-Caucasus*. Moscow (Gost. Ord. Tr. Krasn. Znam. Neft Inst. Gostoptehizdat) (in Russian).
- Sveier, A.V. 1949. Basic morphology and systematics of Pliocene and Post-Pliocene ostracods. *Trudy Vsesoyuznogo Neftyanago Nauchno-Issledovatel'skogo Geologo-Razvedochjnogo Instituta (VNIGRI), Novaya Seriya, Leningrad & Moscow*, **30**: 1–106 (in Russian).
- Szczecchura, J. 1985. *Bolboforma* and similar problematica from the Middle Miocene of Central Paratethys. *Abstracts, VII Congress Regional Commission on Mediterranean Neogene Stratigraphy (RCMNS)*, Budapest, 1985: 541.
- Taner, G. 1985. Die Moluskenfauna der Neogenen Schichten der Anatolischen Seite von Dardanellen. *Abstracts, VII Congress Regional Commission on Mediterranean Neogene Stratigraphy (RCMNS)*, Budapest, 1985: 550–552.
- Tchoumachenko, P., Cernjavskaya, S., Goranov, A., Kojumdzieva, E., Nachev, I., Tikolov, T., Ruskova, N., Sapunov, I. & Tronkov, D. 1990. Principal paleogeographic features of Bulgaria. In: Rakus, M., Dercourt, J. & Nairn, A.E.M. (editors), *Evolution of the Northern Margin of Tethys*. Vol. III: 101–121. Paris.
- Traverse, A. 1978. Palynological analysis of DSDP Leg 42B (1975) Cores from the Black Sea. *Initial Reports of the Deep Sea Drilling Project*, **42B**: 993–1015.
- Trubikhin, V.M. 1977. *Paleomagnetism and stratigraphy of Akchagyljan deposits of west Turkmenia*. Moscow ('Nauka') (in Russian).
- , Bagin, V.I., Gandler, T.S., Nechaeva, T.B. & Fein, A.G. 1991. Late Holocene of west Turkmenia: paleomagnetism, chronology and climatic events. *Abstracts, XIII INQUA International Congress, Beijing, China, 1991*: 359.
- , Nechaeva, T.B. & Fein, A.G. 1991. Ponto-Caspian Quaternary deposits: stratigraphy, chronology, fine structure of magnetic field. *Abstracts, XIII INQUA International Congress, Beijing, China, 1991*: 359.
- Tufescu, M. 1968. *Ammonia tepida* (Cushman) (Ord. Foraminifera): some features

